

A Catalog of Extended Green Objects (EGOs) in the GLIMPSE Survey: A new sample of massive young stellar object outflow candidates

C.J. Cyganowski¹, B.A. Whitney², E. Holden¹, E. Braden¹, C.L. Brogan³, E. Churchwell¹, R. Indebetouw^{3,4}, D. F. Watson⁵, B. L. Babler¹, R. Benjamin⁶, M. Gomez⁷, M. R. Meade¹, M. S. Povich¹, T. P. Robitaille⁸, C. Watson⁹

ccyganow@astro.wisc.edu

ABSTRACT

Using images from the *Spitzer* GLIMPSE Legacy survey, we have identified more than 300 extended 4.5 μm sources (abbreviated EGO, Extended Green Object, for the common coding of the [4.5] band as green in 3-color composite IRAC images). We present a catalog of these EGOs, including integrated flux density measurements at 3.6, 4.5, 5.8, 8.0, and 24 μm from the GLIMPSE and MIPS GAL surveys. The average angular separation between a source in our sample and the nearest IRAS point source is $>1'$. The majority of EGOs are associated with infrared dark clouds (IRDCs), and where high-resolution 6.7 GHz CH_3OH maser surveys overlap the GLIMPSE coverage, EGOs and 6.7 GHz CH_3OH masers are strongly correlated. Extended 4.5 μm emission is thought to trace shocked molecular gas in protostellar outflows; the association of EGOs with IRDCs and 6.7 GHz CH_3OH masers suggests that extended 4.5 μm emission may

¹University of Wisconsin, Madison, WI 53706

²Space Science Institute, 4750 Walnut St. Suite 205, Boulder, CO 80301, bwhitney@spacescience.org

³NRAO, 520 Edgemont Rd, Charlottesville, VA 22903

⁴University of Virginia, Astronomy Dept., P.O. Box 3818, Charlottesville, VA, 22903-0818

⁵Department of Physics and Astronomy, Vanderbilt University, Nashville, TN 37235

⁶Department of Physics, University of Wisconsin at Whitewater, 800 West Main Street, Whitewater, WI 53190

⁷Observatorio Astronomico, Universidad Nacional de Cordoba, Argentina, Laprida 854, 5000 Cordoba, Argentina

⁸SUPA, School of Physics and Astronomy, University of St. Andrews, North Haugh, St. Andrews, UK

⁹Department of Physics, Manchester College, North Manchester, IN 46962

pinpoint outflows specifically from massive protostars. The mid-infrared colors of EGOs lie in regions of color-color space occupied by young protostars still embedded in infalling envelopes.

Subject headings: stars: formation — infrared: stars — ISM: jets and outflows—
Spitzer

1. Introduction

A fundamental limitation in the study of high-mass star formation is the identification of massive young stellar objects (MYSOs), particularly young MYSOs that have not yet formed hypercompact (HC) or ultracompact (UC) H II regions.¹ The largest samples assembled to date have been drawn from the *IRAS* Point Source Catalog (e.g. Sridharan et al. 2002; Molinari et al. 1996), and are limited by its poor resolution (which often blends emission from multiple sources) and positional uncertainty. The *MSX* Point Source Catalog, the basis for the RMS sample (e.g. Urquhart et al. 2008; Hoare et al. 2005, 2004), represents a vast improvement, but emission from clustered sources may still be blended at the resolution of *MSX*.

Recent large-scale, high-angular resolution *Spitzer* surveys of the Galactic plane using the InfraRed Array Camera (IRAC, 3.6, 4.5, 5.8, and 8.0 μm , resolution $<2''$ in all bands; Fazio et al. 2004) and Multiband Imaging Photometer for Spitzer (MIPS, resolution $\sim 6''$ at 24 μm ; Rieke et al. 2004) present an opportunity to compile a new, less-confused sample of IR-selected MYSO candidates. The IRAC 4.5 μm band provides a promising new diagnostic in the search for MYSO candidates. Extended 4.5 μm emission is a conspicuous and ubiquitous feature of *Spitzer* images of known massive star-forming regions (MSFRs), including DR21 (Davis et al. 2007; Smith et al. 2006, and references therein), S255N (Cyganowski et al. 2007), NGC6334I(N) (Hunter et al. 2006), G34.4+0.23 (Shepherd et al. 2007), and IRAS 18566+0408 (Araya et al. 2007). The IRAC 4.5 μm band contains both H₂ ($v=0-0$, S(9,10,11)) lines and CO ($v=1-0$) bandheads (c.f. Figure 1 of Reach et al. 2006). All of these lines may be excited by shocks, such as those expected when protostellar outflows crash into the ambient ISM. Models of protostellar jets by Smith & Rosen (2005) predict that, of the IRAC bands, the integrated H₂ line luminosity will be greatest in band 2 (4.5 μm). Observationally, the extended 4.5 μm emission of the DR21 outflow has been

¹We define MYSOs to be young stellar objects (YSOs) that will eventually become main-sequence O or early B type stars ($M_* \gtrsim 8 M_\odot$).

interpreted as tracing shocked H_2 based on comparisons with NIR H_2 narrowband images and *ISO* spectra (Davis et al. 2007; Smith et al. 2006, and references therein).

Extended $4.5\ \mu\text{m}$ emission is of particular interest as a new MYSO diagnostic because the selection effects are orthogonal to those of traditional methods of identifying MYSO candidates (e.g. applying color criteria to point source catalogs, as done for *IRAS* by Sridharan et al. (2002) and Molinari et al. (1996) and for *MSX* by Hoare et al. (2005, 2004)). Even at the resolution of IRAC, the environments in which MYSOs form—clusters, infrared dark clouds, and nebulous regions—can complicate point-source-based selection methods. Identifying sources with extended $4.5\ \mu\text{m}$ emission targets a population with ongoing outflow activity and active, rapid accretion—a short-lived, and relatively poorly understood, stage of massive protostellar evolution (c.f Zinnecker & Yorke 2007).

We have compiled a catalog of over 300 extended $4.5\ \mu\text{m}$ sources in the GLIMPSE survey, abbreviated EGO (Extended Green Object, for the common coding of the $[4.5]$ band as green in 3-color composite IRAC images). We present this catalog, split into four sub-catalogs (see §3). We present integrated flux densities in the IRAC $4.5\ \mu\text{m}$ band for all catalogued EGOs, and investigate the nature of these sources by examining their correlations with infrared dark clouds (IRDCs) and 6.7 GHz Class II CH_3OH masers. For those EGOs for which counterparts could be identified and photometered in all four IRAC bands and the MIPS $24\ \mu\text{m}$ band, we present five-band photometry (integrated flux densities at $3.6\ \mu\text{m}$, $4.5\ \mu\text{m}$, $5.8\ \mu\text{m}$, $8.0\ \mu\text{m}$, and $24\ \mu\text{m}$), and investigate the EGOs’ colors. In §2 we describe the data and techniques used to compile the catalog, in §3 we present the catalog, and in §4 we discuss the nature of EGOs.

2. The Data

2.1. Identification of EGOs

This paper catalogs EGOs identified in the GLIMPSE survey area ($10^\circ < l < 65^\circ$ and $295^\circ < b < 350^\circ$, $b = \pm 1^\circ$). To identify candidate outflows, we displayed the GLIMPSE $l \times b = 3^\circ \times 2^\circ$ ($1''.2$ per pixel) mosaics² as 3-color images with the IRAC $3.6\ \mu\text{m}$ image in blue, $4.5\ \mu\text{m}$ in green, and $8.0\ \mu\text{m}$ in red. Each image was searched by visual examination for extended green emission, indicating excess at $4.5\ \mu\text{m}$. The source list compiled from this search was then reviewed by two additional authors independently, and only candidates agreed upon by all three authors are included in the catalog. Images in each of the IRAC

²available at http://data.spitzer.caltech.edu/popular/glimpse/20070416_enhanced_v2/3.1x2.4_mosaics/

bands (3.6, 4.5, 5.8, and 8.0 μm), in the MIPS 24 μm band, and a three-color composite (3.6 μm in blue, 4.5 μm in green, and 8.0 μm in red) for all catalogued EGOs are available online; an example is shown in Figure 1.

Most catalogued EGOs are bright and relatively compact, though more extended than point sources (angular extents range from a few to $>30''$). EGOs are readily distinguishable by eye from supernovae remnants (SNR). The IRAC emission from SNR may be dominated by shock-excited warm molecular hydrogen (Neufeld & Yuan 2008), but SNR are generally larger, more diffuse, and more structured than EGOs (Reach et al. 2006). EGOs are also distinguishable from Planetary Nebulae (PNe), which appear red in these 3-color images and generally exhibit spherical morphology at MIR wavelengths. There are three regimes in which our EGO catalog is likely to be incomplete: (1) extended green emission near bright sources; (2) extended green emission in sources with significant PAH emission; and (3) diffuse green emission extended $\lesssim 10''$. Image artifacts associated with bright IRAC sources may obscure or confuse extended 4.5 μm emission associated with these sources. Bright PAH emission (e.g. in a YSO envelope) may overwhelm extended 4.5 μm emission from an outflow. Slightly extended ($\lesssim 10''$) 4.5 μm emission may be confused with multiple nearby or blended point sources (and vice versa); requiring agreement among three observers may exclude some slightly extended sources.³ This paper does not catalog *point-like* sources that appear green in 3-color images for two reasons: (1) point sources with distinctive [3.6]-[4.5] colors may be identified via color selection in the GLIMPSE point source catalog and archive (e.g. Ellingsen 2006, 2007); and (2) due to the flat extinction curve between 4.5 and 8.0 μm (Indebetouw et al. 2005), highly extincted stars could potentially appear green in 3.6/4.5/8.0 μm color images (e.g. Rathborne et al. 2005).

2.2. Photometry

Areal photometry was performed on all catalogued EGOs, using ds9 funtools⁴. Arbitrarily shaped polygon apertures were made to closely match the source morphology in 4.5 μm images. The same polygonal aperture was used for flux density measurements in all four IRAC bands.

A separate aperture was made to measure the local background, the flux density was integrated over the source, and the area-corrected background subtracted. The flux densities

³Of 326 sources identified in the initial search, 24 ($\sim 7\%$) were excluded because the three observers did not agree on their inclusion.

⁴Available at <http://hea-www.harvard.edu/saord/funtools/ds9.html>

of EGOs at $24\ \mu\text{m}$ were also measured, since this wavelength, combined with IRAC, is a good indicator of evolutionary stage in protostars (Robitaille et al. 2006). We measured $24\ \mu\text{m}$ flux densities from mosaics of the Basic Calibrated Data (BCD) from the MIPS GAL survey. Larger, circular or elliptical apertures were used for the $24\ \mu\text{m}$ photometry because the beamsize at $24\ \mu\text{m}$ is at least three times larger than at IRAC wavelengths.

Most EGOs are located in areas of high and/or variable background emission, and uncertainty in the determination of the background level is a dominant contributor to the overall uncertainty in the measured flux densities. For each band, three measurements were made of the source integrated flux density, using three different background regions. Care was taken in placing the background regions to avoid bright stars and areas of exceptionally strong or weak emission. The average of the three measured source flux densities is reported in Tables 1-5, along with the standard deviation of the three measurements. The standard deviation provides an estimate of the uncertainty in the reported source flux densities resulting from uncertain background subtraction. The uncertainty due to background subtraction is generally greatest for the $8\ \mu\text{m}$ flux densities, as diffuse background emission is strongest and most spatially variable in this band. The other major source of uncertainty in the measured source flux densities is observer-dependent aperture selection. To quantify this effect, a sample of 15 sources was photometered by two additional authors; Figure 2 shows the IRAC and MIPS apertures selected by the three different observers for the example source shown in Figure 1. The mean observer-dependent uncertainty is estimated to be $\sim 15\%$ in the IRAC 3.6, 4.5, and $5.8\ \mu\text{m}$ bands and the MIPS $24\ \mu\text{m}$ band and about $\sim 25\%$ in the IRAC $8.0\ \mu\text{m}$ band, in addition to the background-dependent uncertainties listed in Tables 1-5.

3. The Catalogs

Tables 1-4 present the four subcatalogs of EGOs: “likely” MYSO outflow candidates are listed in Tables 1-2, “possible” MYSO outflow candidates in Tables 3-4. Table 1 presents integrated [3.6], [4.5], [5.8], [8.0], and [24] flux densities for “likely” MYSO outflow candidates with counterparts in all five bands (97 sources); for “likely” sources in which confusion and/or saturation are problematic in some bands, Table 2 presents integrated [4.5] flux densities only (36 sources). Similarly, Table 3 presents integrated [3.6], [4.5], [5.8], [8.0], and [24] flux densities for “possible” MYSO outflow candidates with counterparts in all five bands (75 sources); for “possible” sources in which confusion and/or saturation are problematic in some bands, Table 4 presents integrated [4.5] flux densities only (90 sources). Table 5 reports integrated [3.6] and [4.5] flux densities, and [3.6-4.5] colors, for sources in which extended outflow emission may be photometered independently of a central source (see discussion in

§4.3).

The sources in Tables 1-5 are organized by increasing galactic longitude. The columns in Tables 1 and 3 are: (1) name (galactic coordinates), (2-3) position (RA and Dec, rounded to the nearest 0.1 second and nearest arcsecond respectively), (4) the area over which IRAC flux densities were integrated, in units of square arcseconds, (5-8) integrated flux densities in the 3.6, 4.5, 5.8, and 8.0 μm IRAC bands (mJy), (9) the area over which the MIPS 24 μm flux density was integrated, in units of square arcseconds, (10) integrated flux density in the 24 μm MIPS band (mJy), (11) a flag for whether the source is saturated in any band and (12) a list of bands in which the quoted flux density should be considered an upper limit because the measurement may include flux from the psf of a nearby source. Tables 2 and 4 are similar in format, except that only [4.5] flux densities are reported.

The coordinates in columns 2-3 in Tables 1-5 are those of the brightest 4.5 μm emission associated with the EGO. We emphasize that the EGOs in Tables 1-5 have angular extents on the sky ranging from a few to $>30''$, and many are near other MIR GLIMPSE and/or MIPS sources. To establish correlations (or the lack thereof) between the EGOs presented here and other datasets, or to plan followup observations, visual examination of the GLIMPSE and MIPS GAL images should be used along with the listed positions.

The errors quoted in Tables 1-5 are the standard deviations of the three measurements made with different background regions for each band (see §2.2). All reported flux densities are rounded to the nearest 0.1 mJy, and the reported areas are rounded to the nearest square arcsecond.

The categorization of an EGO as a “likely” or “possible” MYSO outflow candidate was based primarily on the angular extent of the extended excess 4.5 μm emission, e.g. the extent of diffuse green emission in 3-color images. Three-color images of each EGO were reviewed by two of the authors; any source in which either observer thought it was possible that multiple nearby point sources and/or image artifacts from a bright IRAC source could be confused with truly extended 4.5 μm emission is considered a “possible” candidate. These relatively compact sources are likely still good YSO candidates (see §4), but not necessarily MYSOs *with outflows* and so likely to be actively accreting.

Even at the resolution of IRAC, the star-forming regions in which some EGOs are located are confused, leading to uncertainty in identifying counterparts in all five mid-infrared (MIR) bands (3.6, 4.5, 5.8, 8.0, and 24 μm) and in placing photometric apertures. Since the spatial resolution of MIPS at 24 μm ($\sim 6''$) is about three times poorer than that of IRAC, sources that are distinct in IRAC images may be blended in MIPS 24 μm images. For each EGO, IRAC 3.6, 4.5, 5.8, and 8.0 μm images and the MIPS 24 μm image were displayed

with the regions chosen for IRAC and MIPS photometry overlaid, as shown in Figure 1a-e. Equivalent figures for all catalogued EGOs are available online. If counterparts to the EGO are distinguishable in all bands (as in Figure 1), then five flux densities are reported (e.g. Tables 1 and 3). Examples of EGOs assigned to Tables 1-4 are shown in, respectively, Figure 1 (Table 1), Figures 3-4 (Table 2), Figure 5 (Table 3), and Figure 6 (Table 4).

4. Discussion: The Nature of EGOs

4.1. Infrared Associations

The only data available for many of the EGOs in Tables 1-5 come from IR all-sky surveys, with *Spitzer* and with previous satellites. To assess whether EGOs constitute a previously unrecognized population of MYSO candidates, Table 6 lists the name and angular separation of the nearest IRAS point source (catalog version 2.1) for each catalogued EGO. Figure 7 graphically presents the data in Table 6. While some EGOs may be identified with IRAS point sources, and a few are well-known MYSOs (G12.91-0.26 (W33a), for example), the mean angular separation between a catalogued EGO and the nearest IRAS point source is $\sim 85''$. The mean angular separation is $\sim 75''$, $\sim 84''$, $\sim 116''$, $\sim 73''$, and $\sim 40''$ for EGOs in Tables 1, 2, 3, 4 and 5, respectively. Figure 8 shows the EGO G11.92-0.61 (shown in Figures 1-2) as seen by *IRAS*, *MSX*, and IRAC. As illustrated by Figure 8, the fact that most EGOs are not *IRAS* point sources does not indicate a lack of emission at the wavelengths of the *IRAS* bands, but rather in many cases is attributable to confusion and/or positional uncertainty at the resolution of *IRAS*.

Infrared dark clouds (IRDCs, c.f. Simon et al. 2006a,b) have been shown in recent years to be sites of the earliest stages of high mass star formation (e.g. Rathborne et al. 2007; Rathborne et al. 2006, and references therein). IRDCs are readily visible against the diffuse background emission in GLIMPSE images, particularly at $8.0 \mu\text{m}$. Many of these are also visible in *MSX* $8.3 \mu\text{m}$ (Band A) images and are catalogued in Simon et al. (2006a) and Simon et al. (2006b). Small and/or filamentary IRDCs, however, may not be evident at the $20''$ resolution of *MSX* Band A (~ 10 times poorer than that of *Spitzer* at $8 \mu\text{m}$). To take advantage of the higher resolution afforded by *Spitzer*, we have analyzed the correlation of EGOs with IRDCs based on visual examination of $8.0 \mu\text{m}$ GLIMPSE images.

The environment of each EGO included in Tables 1-5 was examined in $8.0 \mu\text{m}$ GLIMPSE images and IRDCs identified by visual examination; the results are tabulated in Table 6. (Note that the 3-color IRAC images of each EGO available online are not, in general, scaled to highlight the presence or morphology of dark clouds.) Overall, 67% of the EGOs in Tables 1-

5 are in IRDCs, many of which are small or filamentary. Of our “likely” MYSO outflow candidate EGOs, 71% are associated with IRDCs: 72% in Table 1 and 69% in Table 2. Of our “possible” MYSO outflow candidate EGOs, 64% are associated with IRDCs: 77% in Table 3 and 54% in Table 4. IRDCs are only visible in areas of high diffuse galactic background emission, which generally decreases away from the Galactic Plane. Figure 9 shows the locations of EGOs overplotted on integrated CO emission from Dame et al. (2001), with EGOs from each of Tables 1-5 represented by a different symbol. The overall distribution pattern of EGOs with respect to CO is the same for sources from all subcatalogs: the vast majority of EGOs coincide with CO clouds. For a subset of EGOs for which we had ancillary data, we attempted to determine v_{LSR} from the ^{13}CO Galactic Ring Survey (GRS: Jackson et al. 2006). Due to the presence of multiple ^{13}CO components along the line of sight and the small angular size of EGOs, however, the success rate was only $\sim 50\%$, which was not sufficiently successful to attempt for the whole sample.

The association of extended $4.5\ \mu\text{m}$ emission with known MYSOs (§1), along with the strong correlation of EGOs with IRDCs, suggests that extended $4.5\ \mu\text{m}$ emission in GLIMPSE images may be a good tracer of outflows specifically from massive YSOs. Extended $4.5\ \mu\text{m}$ emission can also originate from low mass outflows (e.g. HH46/47, Noriega-Crespo et al. 2004; Velusamy, Langer, & Marsh 2007). GLIMPSE is, however, a shallow survey: the faintest EGOs identified in GLIMPSE and catalogued in Tables 1-5 have surface brightnesses in $4.5\ \mu\text{m}$ images of $\gtrsim 4\ \text{MJy sr}^{-1}$. In comparison, the bright knots in the HH46/47 outflow have $4.5\ \mu\text{m}$ surface brightness of $\lesssim 2\ \text{MJy sr}^{-1}$, and the majority of the diffuse green emission in NGC1333 (Gutermuth et al. 2008) has $4.5\ \mu\text{m}$ surface brightness of $\lesssim 4\ \text{MJy sr}^{-1}$, with a few very bright areas of $\lesssim 10\ \text{MJy sr}^{-1}$. In both regions, the morphology of the diffuse $4.5\ \mu\text{m}$ emission is mirrored in the other IRAC bands (e.g. Fig.1 of Velusamy, Langer, & Marsh 2007), in contrast to most of the EGOs in our sample (see §4.3). Relatively faint diffuse $4.5\ \mu\text{m}$ emission such as that seen in low-mass star-forming regions is unlikely to have been identified in our search of the GLIMPSE images.

4.2. Association with 6.7 GHz CH_3OH masers

Further evidence that GLIMPSE-identified EGOs are massive comes from the association of EGOs with 6.7 GHz Class II CH_3OH masers. Class II 6.7 GHz CH_3OH masers are indicative of massive star formation (Szymczak, Pillai, & Menten 2005; Ellingsen 2006), as they are *not* observed towards low-mass YSOs (Minier et al. 2003). These masers are radiatively pumped by IR photons emitted by warm dust in massive star-forming regions (e.g. Cragg et al. 2005, and references therein), and are thought to trace both the inner parts

of outflows and disks (c.f. De Buizer 2003). We have compared our EGO catalog with 6.7 GHz Class II CH₃OH maser catalogs of Ellingsen (2006), Walsh et al. (1998), and Caswell (1996). To meaningfully assess the correlation of EGOs and 6.7 GHz CH₃OH masers requires (a) maser positions known to $\sim 1''$ or better, and (b) surveys covering well-defined areas on the sky. To our knowledge, the three surveys cited (and included in the analysis of GLIMPSE point sources associated with 6.7 GHz CH₃OH masers conducted by Ellingsen 2006) are the only studies meeting these criteria that overlap the GLIMPSE coverage. Of these studies, those of Ellingsen (2006) and Caswell (1996) cover (overlapping) ranges in galactic longitude. That of Walsh et al. (1998) targeted *IRAS* sources; some EGOs serendipitously fall within the (large) primary beam. Table 7 tabulates every EGO that falls within the nominal coverage of one of these three surveys (25 EGOs from Table 1, 23 from Table 2, 26 from Table 3, 35 from Table 4, and 2 from Table 5). Column (1) lists the EGO name (galactic coordinates), (2) the maser survey(s) that cover the EGO position, (3) the *IRAS* point source targeted by Walsh et al. (1998) (if relevant), (4) whether a maser was detected, (5) the nominal offset of the maser position from the EGO position (from Tables 1-5), and (6) the distance. The association of 6.7 GHz CH₃OH maser(s) with an EGO was evaluated by plotting maser positions on images in each IRAC and the MIPS 24 μm band and on three-color IRAC images of EGOs, as illustrated in Figure 10a-f. Only masers spatially coincident with either (a) extended 4.5 μm emission, or (b) a MIPS 24 μm source or IRAC multiband source likely to be associated with extended 4.5 μm emission, are tabulated in Table 7. In the latter case, a “?” in column 4 indicates a possible association. Figure 11 graphically presents the nominal offsets between EGO positions (as listed in Tables 1-5) and the maser positions from the literature. As shown in Figure 11, for the vast majority of EGOs this offset is small ($\lesssim 5''$). We emphasize that EGOs are extended on the sky, and a maser with a nominal positional offset of a few arcseconds generally lies within the area of extended 4.5 μm emission. Figures equivalent to Figure 10a-f are available online for all the sources listed in Table 7 to illustrate the range of projected locations of 6.7 GHz CH₃OH masers relative to extended 4.5 μm emission. Among the EGOs in Table 7 are examples of masers coincident with multiband “central” sources, with 4.5 μm emission without clear counterparts in other MIR bands (e.g. the EGO shown in Figures 3 and 10), and with bright MIPS 24 μm point sources slightly offset from extended 4.5 μm emission (e.g. G344.23-0.57).

The non-detections in Table 7 should be treated with caution: all of the maser surveys cited are flux-limited, and the sensitivity of the Walsh et al. (1998) and Caswell (1996) surveys is not uniform, but depends on the position of a source within the interferometric primary beam. This is reflected in the citations for maser detections in column 4 of Table 7; in several cases in which an EGO fell within the nominal coverage of multiple maser surveys, a maser was detected in one survey but not in another. The fraction of EGOs within the

maser survey areas found to be associated with 6.7 GHz CH₃OH masers should thus be regarded as a lower limit: 73% of “likely” MYSO outflow candidate EGOs (61% in Table 1 and 86% in Table 2) and 27% of “possible” MYSO outflow candidate EGOs (4% in Table 3 and 44% in Table 4). Both of the EGOs in Table 5 that fall within the surveys’ coverage have associated 6.7 GHz CH₃OH masers.

Ellingsen (2006) reports kinematic distances for detected 6.7 GHz CH₃OH masers; the near kinematic distances for masers associated with EGOs from Ellingsen (2006) are listed in column 5 of Table 7. The association of EGOs with IRDCs supports the adoption of the near kinematic distance. Where two distances are listed, two masers in the Ellingsen (2006) sample appear to be associated with the EGO. The range of distances—from 2.6 to 5.3 kpc—is typical of distances to MSFR in the Galactic Plane and to IRDCs as determined by Simon et al. (2006b) from ¹³CO emission.

4.3. MIR Colors

As illustrated in Figure 12, the integrated flux densities reported in Tables 1 and 3 are likely dominated by emission from the central source of an EGO, at least at wavelengths $>4.5 \mu\text{m}$. As a result, the integrated flux densities may be used, with caution, to infer the properties of the sources responsible for driving the hypothesized outflows traced by extended $4.5 \mu\text{m}$ emission. Figure 13 shows that EGOs are found in the region of color-color space occupied by very young YSOs, surrounded by substantial accreting envelopes (Robitaille et al. 2006). Detailed spectral energy distribution (SED) fitting has not been attempted because without long-wavelength information the SEDs are poorly constrained, and the resolution of MIPS GAL at $70 \mu\text{m}$ is too poor to resolve EGOs in crowded regions.

The EGO shown in Figure 12 is a rare example of an EGO in which a central source is sufficiently distinct that extended “outflow” emission may be photometered separately. As shown in Figure 12a-e, the extended $4.5 \mu\text{m}$ emission evident in Figure 12b only has a comparably extended counterpart in the $3.6 \mu\text{m}$ band. Table 5 reports $3.6 \mu\text{m}$ and $4.5 \mu\text{m}$ integrated flux densities for the “outflow-only” regions shown in Figure 12f and for four additional sources in which “outflow-only” regions can be distinguished (shown in online Figure 12). The [3.6]-[4.5] colors of the extended emission in these five sources fall within the range determined for “the shocked outflow nebulosity” in the DR21 outflow by Smith et al. (2006) and redward in [3.6]-[4.5] space of the non-outflow “surrounding nebulosity” (c.f. Figure 7 of Smith et al. 2006).

5. Conclusions

We have catalogued over 300 EGOs, identified by their extended $4.5\ \mu\text{m}$ emission, in GLIMPSE images. The suggestion that extended $4.5\ \mu\text{m}$ emission in GLIMPSE traces outflows specifically from *massive* YSOs is supported by the strong correlation of EGOs with IRDCs and 6.7 GHz Class II CH_3OH masers, both associated with early stages of massive star formation. The mid-infrared colors of EGOs lie in regions of color-color space occupied by young protostars still embedded in infalling envelopes. Taken together, this evidence suggests that *Spitzer* images (e.g. the GLIMPSE I, II and 3D surveys and other Galactic programs) may be used to identify a large number of new MYSO candidates *that are actively accreting*, a crucial category of sources for constraining models of high-mass star formation.

The catalog presented here represents a source list for further study. Most of the candidates have not been previously studied, and broadband *Spitzer* images are insufficient to ensure a homogeneous pool of candidates. We have selected a sample of ~ 30 northern EGOs for extensive followup observations designed to test the hypothesis that $4.5\ \mu\text{m}$ emission traces shocked gas in outflows from massive YSOs. This sample was chosen to span a range of $4.5\ \mu\text{m}$ morphologies and associations (or lack thereof) with other tracers of massive star formation. We will report the results of these studies in subsequent papers.

This work is based on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. This work was based on data taken by the GLIMPSE and MIPS GAL *Spitzer Space Telescope* Legacy Programs. This research has made use of NASA’s Astrophysics Data System Bibliographic Services and the SIMBAD database operated at CDS, Strasbourg, France. Support for this work was provided by NASA/JPL through contract 1289524. C.J.C. is supported by a National Science Foundation Graduate Research Fellowship.

REFERENCES

- Araya, E., et al. 2007, ApJ, 669, 1050
- Benjamin, R.A., et al. 2003, PASP, 115, 953
- Caswell, J.L. 1996, MNRAS, 279, 79
- Cragg, D. M., Sobolev, A. M., & Godfrey, P. D. 2005, MNRAS, 360, 533

- Cyganowski, C. J., Brogan, C. L., & Hunter, T. R. 2007, *AJ*, 134, 346
- Dame, T. M., Hartmann, D., & Thaddeus, P. 2001, *ApJ*, 547, 792
- Davis, C.J., Kumar, M.S.N., Sandell, G., Froebrich, D., Smith, M.D., & Currie, M.J. 2007, *MNRAS*, 374, 29
- De Buizer, J. M. 2003, *MNRAS*, 341, 277
- Ellingsen, S. P., von Bibra, M. L., McCulloch, P. M., Norris, R. P., Deshpande, A. A., & Phillips, C. J. 1996, *MNRAS*, 280, 378
- Ellingsen, S.P. 2006, *ApJ*, 638, 241
- Ellingsen, S. P. 2007, *MNRAS*, 377, 571
- Fazio et al. 2004, *ApJS*, 154, 10
- Gutermuth, R. A., et al. 2008, *ApJ*, 674, 336
- Hoare, M.G., Lumsden, S.L., Oudmaijer, R.D., Urquhart, J.S., Busfield, A.L., Sheret, T.L., Clarke, A.J., Moore, T.J.T., Allsopp, J., Burton, M.G., Purcell, C.R., Jiang, Z., & Wang, M. 2005, *IAUS*, 227, 370
- Hoare, M.G., Lumsden, S.L., Oudmaijer, R.D., Busfield, A.L., King, T.L., & Moore, T.L.J. 2004, *ASPC*, 317, 156
- Hunter, T.R., Brogan, C.L., Megeath, S.T., Menten, K.M., Beuther, H. & Thorwirth, S. 2006, *ApJ*, 649, 888.
- Indebetouw, R., et al. 2005, *ApJ*, 619, 931
- Jackson, J. M., et al. 2006, *ApJS*, 163, 145
- Megeath, S. T., Wilson, T. L., & Corbin, M. R. 2005, *ApJ*, 622, L141
- Minier, V., Ellingsen, S.P., Norris, R.P., & Booth, R.S. 2003, *A&A*, 403, 1095
- Molinari, S., Brand, J., Cesaroni, R., & Palla, F. 1996, *A&A*, 308, 573
- Neufeld, D., & Yuan, Y. 2008, *ArXiv e-prints*, 801, arXiv:0801.2141
- Noreiga-Crespo, A., et al. 2004, *ApJS*, 154, 352
- Rathborne, J. M., Simon, R., & Jackson, J. M. 2007, *ApJ*, 662, 1082

- Rathborne, J.M., Jackson, J.M., & Simon, R. 2006, *ApJ*, 641, 389
- Rathborne, J.M., Jackson, J.M., Chambers, E.T., Simon, R., Shipman, R., & Frieswijk, W. 2005, *ApJ*, 630, L181
- Reach, W.T., Rho, J., Tappe, A., Pannuti, T.G., Brogan, C.L., Churchwell, E.B., Meade, M.R., Babler, B., Indebetouw, r., & Whitney, B.A. 2006, *AJ*, 131, 1479
- Rieke, G.H. et al. 2004, *ApJS*, 154, 25
- Robitaille, T.P., Whitney, B.A., Indebetouw, R., Wood, K., & Denzmore, P. 2006, *ApJS*, 167, 256
- Shepherd, D. S., et al. 2007, *ApJ*, 669, 464
- Simon, R., Jackson, J. M., Rathborne, J. M., & Chambers, E. T. 2006, *ApJ*, 639, 227
- Simon, R., Rathborne, J. M., Shah, R. Y., Jackson, J. M., & Chambers, E. T. 2006, *ApJ*, 653, 1325
- Smith, H.A., Hora, J.L., Marengo, M., & Pipher, J.L. 2006, *ApJ*, 645, 1264
- Smith, M.D., & Rosen, A. 2005, *MNRAS*, 357, 1370
- Sridharan, T. K., Beuther, H., Schilke, P., Menten, K. M., & Wyrowski, F. 2002, *ApJ*, 566, 931
- Szymczak, M., Pillai, T., & Menten, K.M. 2005, *A&A*, 434, 613
- Urquhart, J. S., Hoare, M. G., Lumsden, S. L., Oudmaijer, R. D., & Moore, T. J. T. 2008, *Astronomical Society of the Pacific Conference Series*, 387, 381
- Velusamy, T., Langer, W.D., & Marsh, K.A. 2007, *ApJ*, 668, L159
- Walsh, A.J., Burton, M.G., Hyland, A.R., & Robinson, G. 1998, *MNRAS*, 301, 640
- Zinnecker, H., & Yorke, H. W. 2007, *ARA&A*, 45, 481

Table 1. MIR Integrated Flux Densities: “likely” MYSO outflow candidates

Name	J2000 Coordinates		area($''$) ²	Integrated flux(mJy)				area($''$) ²	Integrated flux(mJy)	Saturated?	Upper limit?
	α (h m s)	δ (° ' ")	IRAC	[3.6]	[4.5]	[5.8]	[8.0]	MIPS	[24]		
G11.92-0.61	18 13 58.1	-18 54 17	583	90.7(1.7)	334.4(3.6)	355.8(23.7)	197.8(36.6)	962	7331.9 (154.6)	Y, 24	
G12.02-0.21	18 12 40.4	-18 37 11	73	3.9(2.1)	14.1(1.3)	16.6(11.1)	11.3(1.7)	132	113.5 (26.1)	N	
G12.91-0.03	18 13 48.2	-17 45 39	75	4.9(0.1)	42.7(0.7)	36.8(8.4)	24.3(5.4)	213	2168.7 (49.6)	N	8,24
G14.33-0.64	18 18 54.4	-16 47 46	163	12.1(0.8)	53.8(0.7)	123.7(8.3)	186.3(22.8)	346	4918.9 (73.7)	Y, 24	8,24
G14.63-0.58	18 19 15.4	-16 30 07	79	6.3(0.2)	24.2(0.3)	44.3(1.0)	50.9(5.5)	772	1563.4 (105.0)	N	24
G16.61-0.24	18 21 52.7	-14 35 51	170	15.0(0.3)	43.6(2.8)	43.7(6.4)	21.2(13.3)	559	333.3 (34.7)	N	
G18.67+0.03	18 24 53.7	-12 39 20	111	17.8(2.2)	56.1(2.1)	84.7(0.5)	72.0(13.8)	662	3921.3 (60.5)	N	24
G18.89-0.47	18 27 07.9	-12 41 36	85	6.9(2.2)	25.4(1.8)	24.3(21.5)	38.6(20.9)	138	475.8 (74.1)	N	
G19.01-0.03	18 25 44.8	-12 22 46	721	88.7(15.2)	307.2(12.6)	471.5(80.9)	570.1(88.8)	835	3468.4 (287.3)	N	
G19.88-0.53	18 29 14.7	-11 50 23	423	145.6(24.6)	566.2(15.9)	910.6(78.8)	790.8(135.6)	783	5055.1 (14.8)	N	
G22.04+0.22	18 30 34.7	-09 34 47	75	1.7(1.4)	14.0(0.2)	19.3(3.5)	8.8(6.4)	536	3293.5 (193.8)	N	24
G23.01-0.41	18 34 40.2	-09 00 38	272	33.1(2.3)	225.5(3.6)	446.7(32.9)	221.5(30.7)	973	6189.0 (481.4)	Y,24	
G23.96-0.11	18 35 22.3	-08 01 28	184	30.7(1.6)	172.7(3.0)	245.2(12.4)	92.2(3.0)	852	1636.5 (99.1)	N	
G24.00-0.10	18 35 23.5	-07 59 32	37	0.7(0.4)	8.9(0.7)	16.7(1.1)	13.7(3.4)	69	294.0 (1.6)	N	
G24.17-0.02	18 35 25.0	-07 48 15	46	1.9(0.6)	8.9(0.8)	6.9(2.8)	5.8(5.8)	115	114.4 (4.1)	N	
G24.94+0.07	18 36 31.5	-07 04 16	223	22.3(3.7)	55.3(1.6)	68.1(1.7)	49.8(12.1)	818	1383.1 (116.5)	N	
G25.27-0.43	18 38 57.0	-07 00 48	262	14.6(3.2)	43.3(2.4)	32.9(21.7)	16.2(4.4)	156	190.8 (4.2)	N	
G27.97-0.47	18 44 03.6	-04 38 02	202	8.6(1.2)	48.0(0.3)	83.1(12.1)	102.6(30.8)	484	1820.7 (36.4)	N	24
G28.83-0.25	18 44 51.3	-03 45 48	575	52.3(15.2)	171.0(19.5)	115.4(52.9)	142.5(19.5)	478	3635.9 (64.6)	N	8,24
G34.41+0.24	18 53 17.9	+01 25 25	114	7.2(0.4)	21.4(0.4)	24.9(4.3)	16.5(10.7)	864	6227.1 (32.4)	Y,24	
G35.03+0.35	18 54 00.5	+02 01 18	845	161.6(11.8)	501.1(7.3)	511.9(217.8)	279.6(438.5)	1089	10181.9 (231.7)	Y,24	3,4,5,8,24
G35.04-0.47	18 56 58.1	+01 39 37	154	8.5(1.7)	25.4(0.8)	17.6(0.5)	11.8(5.8)	426	188.8 (16.0)	N	3,4,5,8,24
G35.13-0.74	18 58 06.4	+01 37 01	145	97.5(0.8)	181.1(3.0)	327.2(8.0)	452.2(43.7)	104	1305.5 (31.1)	N	8,24
G35.15+0.80	18 52 36.6	+02 20 26	351	97.6(2.0)	133.7(2.3)	275.6(1.4)	367.9(9.4)	714	1880.2 (33.8)	N	3,4,5,8,24
G35.20-0.74	18 58 12.9	+01 40 33	1066	490.3(9.3)	1080.3(2.7)	1623.1(38.2)	2879.6(97.3)	9971	31704.5 (530.1)	Y,24	
G35.68-0.18	18 57 05.0	+02 22 00	117	5.2(0.4)	23.3(0.8)	8.7(0.2)	7.9(2.5)	449	485.6 (40.5)	N	8,24
G35.79-0.17	18 57 16.7	+02 27 56	48	3.2(0.6)	12.2(0.3)	14.5(1.3)	11.4(6.5)	668	1303.1 (39.0)	N	
G36.01-0.20	18 57 45.9	+02 39 05	109	5.4(1.6)	18.0(1.7)	12.1(0.8)	5.8(1.5)	109	59.3 (3.5)	N	
G37.48-0.10	19 00 07.0	+03 59 53	107	7.5(1.2)	19.0(0.8)	40.9(4.0)	60.7(7.1)	714	1096.3 (364.0)	N	
G39.10+0.49	19 00 58.1	+05 42 44	120	11.9(0.3)	29.2(1.0)	23.0(0.9)	9.7(1.8)	876	1439.8 (9.1)	N	
G40.28-0.27	19 05 51.5	+06 24 39	114	17.2(0.2)	73.4(0.4)	127.3(3.4)	107.7(8.1)	2604	3221.2 (87.5)	N	24
G44.01-0.03	19 11 57.2	+09 50 05	117	30.0(0.7)	31.7(0.6)	29.6(1.1)	11.8(5.8)	63	16.0 (3.8)	N	
G45.47+0.05	19 14 25.6	+11 09 28	248	65.0(3.7)	183.3(2.8)	143.7(20.5)	91.5(69.5)	847	15702.4 (806.6)	Y,24	
G49.27-0.34	19 23 06.7	+14 20 13	441	82.5(11.2)	666.3(8.0)	1737.2(42.9)	1880.6(144.9)	1319	8677.1 (174.5)	Y,24	

Table 1—Continued

Name	J2000 Coordinates		area('') ² IRAC	Integrated flux(mJy)				area('') ² MIPS	Integrated flux(mJy) [24]	Saturated?	Upper limit?
	α (h m s)	δ (° ' ")		[3.6]	[4.5]	[5.8]	[8.0]				
G54.45+1.02	19 28 25.7	+19 32 20	50	4.5(0.2)	9.4(0.0)	15.6(1.3)	22.7(1.8)	52	93.2 (2.4)	N	24
G56.13+0.22	19 34 51.5	+20 37 28	62	3.6(0.1)	9.0(0.1)	17.0(0.4)	22.3(0.6)	207	153.5 (2.4)	N	24
G58.09-0.34	19 41 03.9	+22 03 39	79	10.0(0.1)	12.2(0.3)	7.2(0.3)	1.1(0.8)	582	57.4 (1.4)	N	
G59.79+0.63	19 41 03.1	+24 01 15	121	7.8(0.9)	26.9(0.7)	45.8(3.3)	41.9(10.7)	518	417.4 (29.0)	N	
G298.26+0.74	12 11 47.7	-61 46 21	356	317.1(1.6)	605.6(0.3)	648.8(4.5)	536.3(8.0)	876	18226.2 (51.3)	Y,24	24
G298.89+0.37	12 16 37.9	-62 13 41	35	1.9(0.1)	3.4(0.2)	2.7(0.4)	0.6(0.5)	86	20.2 (3.0)	N	
G298.90+0.36	12 16 43.2	-62 14 25	278	71.7(1.6)	125.7(0.7)	160.1(4.6)	162.0(7.9)	2540	3293.1 (211.9)	N	24
G305.48-0.10	13 13 45.8	-62 51 28	376	107.8(4.3)	275.4(0.9)	358.5(16.5)	304.1(35.9)	639	5052.2 (568.7)	Y, 24	
G305.52+0.76	13 13 29.3	-61 59 53	228	20.7(1.0)	48.7(0.1)	55.8(3.1)	22.8(3.8)	783	645.6 (45.5)	N	
G305.57-0.34	13 14 49.1	-63 05 38	141	20.3(0.7)	35.9(0.6)	25.0(7.1)	13.3(2.9)	536	343.9 (15.0)	N	
G305.62-0.34	13 15 11.5	-63 05 30	85	15.4(0.8)	27.1(0.3)	27.2(3.9)	19.0(8.7)	478	1082.8 (25.1)	N	
G305.82-0.11	13 16 48.6	-62 50 35	84	15.6(0.2)	57.5(0.4)	64.6(1.3)	50.0(9.2)	484	1062.2 (185.6)	N	24
G305.89+0.02	13 17 15.5	-62 42 24	124	36.7(0.9)	120.4(0.2)	166.4(14.1)	158.5(19.6)	622	3639.0 (77.5)	N	24
G309.15-0.35	13 45 51.3	-62 33 46	60	1.8(0.1)	7.2(0.2)	6.6(2.9)	8.0(4.9)	81	9.3 (3.3)	N	
G309.38-0.13(a)	13 47 23.9	-62 18 12	131	8.2(0.3)	20.3(1.2)	26.2(4.4)	9.4(9.6)	645	3852.7 (54.8)	N	
G309.99+0.51(a)	13 51 12.2	-61 32 09	95	3.9(1.0)	8.8(0.1)	7.8(2.6)	15.7(6.4)	92	69.3 (1.1)	N	
G310.08-0.23	13 53 23.0	-62 14 13	128	53.1(1.2)	121.6(0.8)	183.6(1.0)	161.7(9.9)	605	2959.2 (95.5)	N	
G312.11+0.26	14 08 49.3	-61 13 25	98	10.2(0.6)	29.0(0.4)	17.3(8.7)	10.2(0.7)	708	2713.6 (44.8)	N	
G317.42-0.67	14 51 59.2	-60 06 06	102	22.5(1.0)	106.3(0.5)	222.7(3.8)	242.0(8.7)	668	1173.9 (8.3)	N	24
G317.46-0.40b	14 51 19.6	-59 50 51	89	10.8(1.6)	55.1(0.4)	103.4(2.4)	117.2(14.9)	893	3933.1 (118.5)	Y,24	24
G317.87-0.15	14 53 16.3	-59 26 36	72	6.4(0.1)	13.1(0.2)	10.5(2.4)	8.0(5.4)	161	192.8 (13.7)	N	
G317.88-0.25	14 53 43.5	-59 31 35	73	11.4(0.7)	22.1(0.1)	19.4(4.5)	19.0(1.1)	46	17.5 (1.0)	N	24
G321.94-0.01	15 19 43.3	-57 18 06	102	18.9(1.1)	98.3(0.6)	190.8(2.6)	173.5(5.9)	570	4416.1 (57.4)	N	24
G324.19+0.41	15 31 38.0	-55 42 36	115	25.9(0.6)	32.2(0.7)	53.0(3.6)	53.7(7.2)	138	207.0 (1.0)	N	8,24?
G324.72+0.34	15 34 57.5	-55 27 26	462	26.4(3.6)	86.5(2.5)	65.6(14.9)	12.8(7.1)	708	896.7 (10.8)	N	24
G326.27-0.49	15 47 10.8	-55 11 12	547	147.4(11.8)	494.5(7.1)	921.6(14.1)	784.5(48.6)	737	3139.6 (173.1)	N	8,24
G326.31+0.90	15 41 35.9	-54 03 42	53	13.5(0.5)	27.2(0.2)	36.0(4.3)	45.3(3.7)	121	68.2 (27.5)	N	
G326.78-0.24	15 48 55.2	-54 40 37	295	207.5(0.7)	614.1(0.5)	1085.6(8.5)	1415.1(18.1)	634	9509.5 (101.5)	Y, 24	
G326.79+0.38	15 46 20.9	-54 10 45	406	171.0(2.9)	305.0(3.2)	348.5(25.0)	289.1(23.0)	697	3762.9 (26.1)	N	
G326.86-0.67	15 51 13.6	-54 58 03	226	13.0(0.9)	35.8(2.1)	62.8(4.9)	13.6(11.2)	109	130.9 (11.2)	N	
G326.97-0.03	15 49 03.2	-54 23 37	46	2.4(0.1)	11.1(0.5)	16.8(1.2)	11.2(3.1)	104	67.3 (7.9)	N	24
G327.12+0.51	15 47 32.7	-53 52 39	1048	365.3(13.0)	964.5(4.3)	2544.5(11.3)	4347.8(47.4)	962	19185.4 (143.3)	Y, 24	5.8,8
G327.39+0.20	15 50 18.5	-53 57 07	170	67.5(0.1)	196.5(0.4)	259.9(4.4)	242.4(24.9)	547	3322.0 (23.5)	N	24
G327.40+0.44	15 49 19.3	-53 45 10	318	235.6(1.7)	876.7(5.3)	1964.7(7.4)	2829.9(10.5)	841	11960.0 (150.3)	Y,24	8,24

- 15 -

Table 1—Continued

Name	J2000 Coordinates		area($''^2$)	Integrated flux(mJy)					area($''^2$)	Integrated flux(mJy)	Saturated?	Upper limit?
	α (h m s)	δ ($^\circ$ ' ")	IRAC	[3.6]	[4.5]	[5.8]	[8.0]	MIPS	[24]			
G328.14-0.43	15 56 57.6	-53 57 48	89	64.9(1.3)	146.9(0.7)	127.8(4.4)	23.8(7.8)	564	2567.3 (15.0)	N		
G329.18-0.31	16 01 47.4	-53 11 44	189	18.2(1.6)	90.0(1.4)	89.7(4.4)	31.7(11.7)	662	3969.5 (28.7)	N		24
G329.47+0.52	15 59 36.6	-52 22 55	117	10.3(0.8)	28.3(1.1)	33.7(2.6)	6.5(3.8)	536	1167.9 (30.8)	N		
G329.61+0.11	16 02 03.1	-52 35 33	274	110.8(0.9)	498.7(0.7)	880.5(21.7)	1431.7(60.6)	3312	8092.8 (438.8)	N		24
G332.35-0.12	16 16 07.0	-50 54 30	308	198.6(3.4)	429.4(1.9)	636.1(26.7)	612.0(70.5)	605	5557.2 (162.1)	Y,24		24
G332.56-0.15	16 17 12.1	-50 47 14	92	20.7(1.6)	51.0(2.2)	34.6(2.5)	21.1(26.5)	115	782.4 (197.7)	N		
G332.81-0.70	16 20 48.1	-51 00 15	160	63.2(4.6)	182.2(5.1)	178.6(14.6)	180.9(81.0)	570	3104.6 (458.5)	N		
G332.94-0.69	16 21 18.9	-50 54 10	240	44.7(3.8)	215.0(2.7)	336.7(36.1)	354.9(43.1)	559	2689.8 (550.2)	N		8
G332.96-0.68	16 21 22.9	-50 52 58	164	34.7(0.6)	47.9(0.8)	283.0(6.0)	705.3(11.8)	444	3622.0 (56.8)	Y,24		8
G333.18-0.09	16 19 45.6	-50 18 34	206	129.2(5.5)	338.1(4.0)	344.5(9.7)	107.0(85.4)	121	1270.5 (137.7)	N		8
G335.59-0.29	16 30 58.5	-48 43 51	982	132.1(11.9)	483.8(9.9)	466.7(54.5)	111.3(134.4)	2160	10230.0 (244.0)	Y,24		
G337.30-0.87	16 40 31.3	-47 51 31	199	56.5(1.3)	111.2(0.7)	103.8(6.8)	38.5(22.3)	547	1356.2 (67.6)	N		
G338.39-0.40	16 42 41.2	-46 43 40	907	175.8(5.4)	336.0(12.4)	300.4(20.1)	117.6(63.5)	553	3298.4 (63.0)	N		8,24
G339.95-0.54	16 49 07.9	-45 37 59	170	41.7(1.1)	140.3(0.8)	188.4(12.5)	231.6(6.9)	766	7877.0 (182.9)	N		
G340.97-1.02	16 54 57.3	-45 09 04	304	12.2(5.7)	95.5(0.1)	75.0(30.2)	44.4(17.8)	680	591.3 (89.5)	N		8
G341.24-0.27	16 52 37.3	-44 28 09	153	20.8(1.2)	57.6(0.1)	60.6(13.9)	38.0(18.5)	536	1189.0 (64.0)	N		8,24
G341.73-0.97	16 57 23.1	-44 31 35	334	73.3(2.8)	213.2(0.8)	295.7(6.2)	230.9(9.1)	530	1184.9 (13.1)	N		
G341.99-0.10	16 54 32.8	-43 46 45	276	48.8(2.7)	162.0(2.3)	192.8(7.6)	89.1(11.7)	691	2304.0 (30.4)	N		
G343.12-0.06	16 58 16.6	-42 52 04	894	196.2(2.6)	875.6(2.7)	1254.9(21.7)	796.6(90.2)	3519	19300.2 (512.4)	Y,24		
G343.50-0.47	17 01 18.4	-42 49 36	204	44.4(0.7)	204.9(1.0)	301.2(8.7)	216.7(8.4)	657	3482.4 (41.8)	N		24
G343.72-0.18(a)	17 00 48.3	-42 28 25	197	33.6(6.6)	60.2(1.3)	55.3(3.5)	32.6(11.7)	495	905.6 (21.7)	N		24
G343.72-0.18(b)	17 00 48.1	-42 28 37	59	4.5(0.6)	15.4(0.1)	18.3(4.4)	31.4(4.9)	472	1092.5 (107.0)	N		24
G344.58-0.02	17 02 57.7	-41 41 54	259	58.0(1.1)	205.8(2.1)	320.0(5.2)	430.1(11.7)	3663	12302.6 (43.5)	Y, 24		8,24
G345.72+0.82	17 03 06.4	-40 17 09	1231	149.2(6.6)	694.6(2.6)	1400.5(29.6)	1371.7(80.6)	3698	7499.2 (73.8)	Y, 24		8,24
G345.99-0.02	17 07 27.6	-40 34 45	76	4.7(0.2)	20.2(1.5)	35.4(1.6)	11.8(1.2)	207	514.3 (8.1)	N		24
G347.08-0.40	17 12 27.2	-39 55 18	72	51.2(0.3)	100.7(0.0)	101.8(2.2)	51.8(6.0)	173	2286.4 (5.5)	N		24
G348.55-0.98	17 19 20.9	-39 03 55	438	198.9(3.4)	442.9(11.4)	650.0(39.7)	680.9(162.8)	564	6324.5 (109.0)	N		8,24
G348.58-0.92	17 19 10.7	-39 00 23	301	75.9(1.7)	216.8(2.8)	246.0(2.9)	154.2(21.2)	576	5098.8 (118.3)	Y, 24		
G349.15-0.98	17 21 04.8	-38 34 25	279	177.9(2.6)	355.6(2.1)	627.0(3.1)	918.5(6.6)	3197	4861.4 (159.7)	Y, 24		24

Table 2. [4.5] Integrated Flux Densities: ”likely” MYSO outflow candidates

Name	J2000 Coordinates		area($''$) ²	Integrated flux(mJy)
	α (h m s)	δ ($^{\circ}$ ' ")	IRAC	[4.5]
G10.29-0.13	18 08 49.3	-20 05 57.3	69	13.1(1.4)
G10.34-0.143	18 09 00.0	-20 03 35	94	72.7(4.3)
G16.59-0.05	18 21 09.1	-14 31 48	128	50.3(0.7)
G19.36-0.03	18 26 25.8	-12 03 56.9	153	115.0(2.1)
G19.61-0.12	18 27 13.6	-11 53 20	102	32.4(5.1)
G25.38-0.15	18 38 08.1	-06 46 53	46	46.4(0.9)
G28.28-0.36	18 44 13.2	-04 18 04	112	47.5(3.3)
G34.39+0.22	18 53 19.0	+01 24 08	307	20.4(4.3)
G48.66-0.30	19 21 48.0	+13 49 21	120	2.0(0.9)
G49.42+0.33	19 20 59.1	+14 46 53	98	12.7(0.9)
G309.90+0.23	13 51 00.4	-61 49 53	109	16.0(0.2)
G317.46-0.40a	14 51 19.6	-59 50 41	39	15.3(0.4)
G318.04+0.09	14 53 39.2	-59 09 10	84	19.6(0.2)
G318.05+0.09	14 53 42.6	-59 08 49	202	419.5(1.3)
G320.23-0.28	15 09 52.6	-58 25 36	170	82.4(0.8)
G324.17+0.44	15 31 24.6	-55 41 30	84	16.4(0.1)
G326.32-0.39	15 47 04.8	-55 04 51	91	19.1(0.0)
G326.47+0.70	15 43 15.4	-54 07 14	69	7.8(0.2)
G326.48+0.70	15 43 17.5	-54 07 11	164	92.5(0.3)
G326.61+0.80(b)	15 43 34.6	-53 58 00	84	17.7(0.6)
G328.25-0.53	15 57 59.7	-53 58 00	523	866.3(3.0)
G329.03-0.20	16 00 30.6	-53 12 34	397	81.6(1.7)
G329.41-0.46	16 03 32.4	-53 09 26	219	258.9(3.1)
G329.47+0.50	15 59 41.0	-52 23 28	219	59.0(1.3)
G330.88-0.37	16 10 19.9	-52 06 13	88	41.9(1.4)
G331.13-0.24	16 10 59.8	-51 50 19	46	21.5(0.9)
G332.12+0.94	16 10 30.4	-50 18 05	121	43.7(0.7)
G332.60-0.17	16 17 29.4	-50 46 13	82	14.2(2.0)
G332.73-0.62	16 20 02.8	-51 00 32	186	80.1(3.2)
G333.47-0.16	16 21 20.2	-50 09 50	117	125.8(0.9)
G335.06-0.43	16 29 23.1	-49 12 28	157	46.5(1.9)
G335.79+0.18	16 29 47.1	-48 15 47	134	93.7(1.9)
G337.91-0.48	16 41 10.3	-47 08 06	613	1981.1(6.0)
G341.22-0.26(a)	16 52 32.2	-44 28 38	158	24.2(4.4)
G342.48+0.18	16 55 02.6	-43 13 01	179	42.6(2.3)
G344.23-0.57	17 04 07.1	-42 18 42	81	18.7(0.5)

Table 3. MIR Integrated Flux Densities: "possible" MYSO outflow candidates

Name	J2000 Coordinates		area('') ²	Integrated flux(mJy)				area('') ²	Integrated flux(mJy)	Saturated?	Upper limit?
	α (h m s)	δ (° ' ")	IRAC	[3.6]	[4.5]	[5.8]	[8.0]	MIPS	[24]		
G11.11-0.11	18 10 28.3	-19 22 31	177	22.2(3.7)	200.7(1.6)	386.7(6.1)	263.1(24.7)	812	1542.6 (80.2)	N	
G16.58-0.08	18 21 15.0	-14 33 02	82	63.7(0.5)	63.5(0.5)	122.6(6.5)	83.1(23.8)	559	497.9 (16.6)	N	24
G24.63+0.15	18 35 40.1	-07 18 35	52	8.4(0.9)	59.8(1.7)	103.1(1.5)	57.9(3.5)	530	405.1 (125.9)	N	24
G29.84-0.47	18 47 28.8	-02 58 03	78	1.4(0.8)	14.0(0.2)	11.2(2.2)	7.4(3.2)	202	36.4 (12.4)	N	
G29.96-0.79	18 48 50.0	-03 00 21	232	82.6(1.5)	128.7(1.5)	227.2(17.4)	367.2(2.4)	167	199.8 (19.9)	N	8
G34.28+0.18	18 53 15.0	+01 17 11	157	13.3(3.2)	47.7(1.5)	68.1(7.3)	44.8(47.0)	444	499.7 (5.7)	N	
G40.28-0.22	19 05 41.3	+06 26 13	225	80.6(0.8)	236.3(0.3)	249.1(0.5)	204.8(1.9)	3214	7733.8 (198.0)	N	24
G45.80-0.36	19 16 31.1	+11 16 11	65	19.1(0.2)	63.0(0.3)	103.8(1.4)	108.1(3.9)	109	854.1 (1.2)	N	24
G49.07-0.33	19 22 41.9	+14 10 12	48	23.3(0.6)	85.9(1.0)	135.3(0.3)	66.1(18.9)	98	381.0 (57.8)	N	
G49.27-0.32	19 23 02.2	+14 20 52	30	2.9(0.7)	5.6(0.7)	3.9(3.5)	6.9(1.6)	524	855.0 (180.1)	N	
G50.36-0.42	19 25 32.8	+15 15 38	82	14.4(0.5)	20.5(0.4)	21.3(0.6)	11.6(2.3)	576	245.4 (0.3)	N	
G53.92-0.07	19 31 23.0	+18 33 00	105	58.9(0.2)	108.0(0.6)	149.9(0.8)	171.5(0.9)	616	1043.1 (45.1)	N	24
G54.11-0.08	19 31 48.8	+18 42 57	58	54.2(0.7)	130.4(0.7)	244.4(5.2)	359.0(15.5)	559	2748.6 (542.0)	N	
G54.45+1.01	19 28 26.4	+19 32 15	91	23.7(0.3)	75.8(0.4)	147.9(1.8)	176.6(6.2)	588	1880.5 (26.4)	N	24
G58.79+0.63	19 38 55.3	+23 09 04	94	7.0(0.1)	12.3(0.1)	13.8(0.5)	16.4(0.6)	230	88.6 (1.6)	N	
G62.70-0.51	19 51 51.1	+25 57 40	99	47.8(0.2)	64.7(0.2)	77.1(0.3)	74.4(1.2)	570	190.0 (3.8)	N	
G304.89+0.64	13 08 12.1	-62 10 22	72	45.7(0.5)	102.9(0.2)	165.8(2.7)	175.1(10.2)	3185	7645.7 (32.7)	Y, 24	
G309.91+0.32	13 50 53.9	-61 44 22	251	61.1(1.1)	129.4(2.1)	96.7(18.5)	20.5(3.2)	467	581.0 (192.8)	N	24
G309.97+0.59	13 50 52.6	-61 27 46	56	2.2(0.5)	4.0(0.1)	3.2(0.7)	3.0(1.7)	81	9.1 (0.9)	N	
G309.99+0.51(b)	13 51 12.7	-61 32 22	49	3.1(0.4)	7.8(0.3)	10.7(2.0)	2.3(1.7)	63	33.2 (2.7)	N	
G311.51-0.45	14 05 46.1	-62 04 49	190	397.1(1.5)	776.0(0.8)	1183.6(12.8)	1507.7(14.4)	3070	10139.1 (1231.5)	Y,24	24
G324.11+0.44	15 31 05.0	-55 43 39	354	58.9(1.3)	185.2(0.7)	207.4(4.8)	171.3(8.3)	737	1537.6 (7.1)	N	
G326.36+0.88	15 41 55.4	-54 02 55	50	2.8(0.8)	9.2(0.5)	12.2(3.2)	13.8(5.1)	104	74.7 (57.1)	N	
G326.41+0.93	15 41 59.4	-53 59 03	343	75.7(3.1)	323.6(3.1)	468.2(24.5)	333.4(19.0)	772	2551.9 (208.0)	N	8
G326.57+0.20	15 45 53.4	-54 27 50	50	4.4(0.4)	17.9(0.1)	25.3(0.8)	16.8(1.2)	132	273.4 (2.5)	N	
G326.61+0.80(a)	15 43 35.1	-53 57 36	150	20.2(0.5)	72.9(0.8)	115.6(4.0)	181.7(19.0)	484	2248.1 (16.9)	N	24
G326.61+0.80(c)	15 43 36.2	-53 57 51	125	92.7(1.4)	162.4(0.5)	174.2(6.1)	150.1(16.0)	501	1816.6 (18.0)	N	24
G326.65+0.75	15 44 00.9	-53 58 45	140	40.5(0.7)	196.8(0.4)	300.3(4.5)	218.2(15.8)	507	2110.8 (27.9)	N	24
G326.80+0.51	15 45 48.6	-54 04 30	81	3.7(0.3)	10.2(0.4)	5.8(3.8)	6.7(3.0)	52	10.0 (2.6)	N	
G326.92-0.31	15 49 56.2	-54 38 29	166	11.0(0.9)	60.6(0.1)	97.8(9.6)	84.3(30.2)	599	551.4 (113.4)	N	24
G327.30-0.58	15 53 11.2	-54 36 48	89	24.5(1.5)	158.8(1.7)	341.8(8.8)	364.9(34.1)	369	2589.1 (86.5)	N	24
G327.72-0.38	15 54 32.3	-54 11 55	37	5.1(0.2)	16.0(0.6)	37.6(3.3)	54.2(6.1)	75	113.0 (19.7)	N	8,24
G327.86+0.19	15 52 49.2	-53 40 07	52	7.6(0.1)	15.9(0.4)	26.2(1.5)	28.8(3.6)	305	339.6 (45.1)	N	8,24
G327.89+0.15	15 53 10.3	-53 40 28	89	10.9(0.5)	38.6(0.2)	57.9(4.0)	43.4(4.9)	179	215.0 (37.9)	N	

Table 3—Continued

Name	J2000 Coordinates		area($''^2$)	Integrated flux(mJy)				area($''^2$)	Integrated flux(mJy)	Saturated?	Upper limit?
	α (h m s)	δ ($^\circ$ ' ")	IRAC	[3.6]	[4.5]	[5.8]	[8.0]	MIPS	[24]		
G328.55+0.27	15 56 01.5	-53 09 44	157	85.1(2.5)	268.8(1.2)	368.4(10.7)	318.4(31.4)	812	4483.8 (57.9)	Y,24	
G329.07-0.31(a)	16 01 11.7	-53 16 00	158	79.8(0.9)	277.9(1.7)	423.4(5.7)	424.4(16.2)	668	4192.3 (39.4)	N	24
G329.07-0.31(b)	16 01 09.9	-53 16 02	217	57.5(0.6)	322.8(0.4)	795.3(5.5)	1054.1(8.7)	795	6590.2 (45.9)	Y,24	8,24
G329.16-0.29	16 01 33.6	-53 11 15	36	4.3(0.3)	11.8(0.0)	16.0(1.1)	11.8(6.3)	75	106.5 (1.1)	N	24
G329.18-0.30	16 01 44.9	-53 11 15	167	15.5(0.8)	66.0(1.3)	104.6(4.1)	50.3(19.3)	645	581.7 (14.6)	N	8
G331.08-0.47	16 11 46.9	-52 02 31	111	6.6(1.0)	26.6(1.7)	60.2(5.2)	58.7(11.9)	415	629.1 (79.4)	N	
G331.37-0.40	16 12 48.1	-51 47 30	86	2.1(1.3)	9.7(0.8)	4.3(3.4)	5.0(1.4)	213	166.9 (11.3)	N	
G331.62+0.53	16 09 56.8	-50 56 25	363	47.6(5.6)	117.2(2.0)	233.2(15.1)	444.2(25.0)	714	2955.7 (75.8)	N	
G331.71+0.58	16 10 06.3	-50 50 29	161	10.7(1.0)	26.5(0.6)	16.6(5.0)	10.6(6.2)	334	245.5 (11.7)	N	
G331.71+0.60	16 10 01.9	-50 49 33	520	122.5(2.2)	315.3(1.4)	421.9(5.5)	395.8(22.8)	876	3720.8 (38.6)	N	
G332.28-0.07	16 15 35.1	-50 55 36	53	9.8(0.3)	29.2(1.9)	24.1(8.8)	14.9(6.1)	150	551.9 (71.1)	N	24
G332.33-0.12	16 16 03.3	-50 55 34	48	8.1(0.9)	20.4(0.1)	26.9(2.5)	23.7(6.6)	98	68.3 (7.8)	N	8
G332.36+0.60	16 13 02.4	-50 22 39	207	102.2(2.1)	237.5(1.2)	368.7(1.6)	382.8(8.4)	714	2216.9 (93.4)	N	
G332.47-0.52	16 18 26.5	-51 07 12	544	149.5(4.6)	420.9(2.8)	1029.5(73.0)	1898.5(105.5)	870	8610.4 (182.3)	Y,24	
G332.58+0.15	16 16 00.6	-50 33 30	69	7.1(0.7)	22.0(0.9)	24.3(4.0)	19.2(6.4)	167	203.9 (17.9)	N	
G332.59+0.04(a)	16 16 30.4	-50 37 41	30	3.4(0.4)	8.8(0.6)	4.8(2.8)	6.0(6.6)	52	22.4 (0.9)	N	
G332.59+0.04(b)	16 16 30.1	-50 37 50	52	7.9(0.6)	20.5(0.5)	27.2(7.2)	41.1(6.5)	52	38.1 (1.9)	N	
G332.91-0.55	16 20 32.6	-50 49 46	75	14.5(0.6)	47.4(0.8)	42.5(5.9)	8.4(8.3)	138	240.2 (70.8)	N	
G333.08-0.56	16 21 20.9	-50 43 05	151	37.7(3.6)	277.3(0.7)	558.8(37.7)	570.9(68.6)	513	1500.1 (187.1)	N	
G334.04+0.35	16 21 36.9	-49 23 28	177	31.0(1.5)	101.9(0.4)	171.8(1.4)	218.7(8.7)	645	2344.9 (18.3)	N	
G334.25+0.07	16 23 45.2	-49 26 32	86	8.8(0.3)	23.4(0.6)	24.1(2.8)	8.7(6.4)	138	201.0 (11.3)	N	
G335.43-0.24	16 30 05.8	-48 48 44	252	38.0(2.3)	298.1(3.5)	481.7(32.1)	266.4(28.8)	570	2380.3 (39.3)	N	
G336.87+0.29	16 33 40.3	-47 23 32	147	21.8(3.2)	66.6(2.5)	62.1(22.5)	39.2(6.2)	478	612.7 (15.1)	N	
G336.96-0.98	16 39 37.5	-48 10 58	125	27.1(0.2)	64.6(0.3)	64.1(7.2)	42.0(2.0)	628	708.6 (67.2)	N	
G337.16-0.39	16 37 49.6	-47 38 50	222	228.3(0.7)	536.9(1.8)	1136.5(15.3)	2237.2(43.8)	1140	5589.0 (144.0)	N	24
G338.32-0.41	16 42 27.5	-46 46 57	193	39.5(1.7)	102.5(1.2)	101.5(6.2)	30.9(5.9)	743	1479.0 (46.8)	N	8,24
G340.75-1.00	16 54 04.0	-45 18 50	320	101.3(6.3)	239.0(1.1)	336.4(28.2)	552.6(14.9)	3554	17376.0 (370.9)	Y,24	24
G340.77-0.12	16 50 17.5	-44 43 54	111	13.1(0.6)	34.3(0.9)	36.3(3.9)	28.6(13.3)	472	358.6 (32.1)	N	
G340.78-0.10	16 50 14.7	-44 42 31	91	6.7(1.0)	22.0(0.6)	25.4(6.2)	20.0(11.3)	726	1737.9 (58.7)	N	24
G341.20-0.26	16 52 27.8	-44 29 29	58	23.3(0.7)	32.3(0.1)	33.8(3.1)	15.5(9.1)	144	49.4 (9.9)	N	
G341.23-0.27	16 52 34.2	-44 28 36	45	6.7(0.3)	14.2(0.2)	18.8(1.1)	13.6(6.0)	115	70.3 (7.3)	N	
G342.15+0.51	16 52 28.3	-43 16 08	50	6.6(0.2)	12.4(0.1)	15.7(0.4)	16.4(0.6)	121	70.5 (2.5)	N	
G342.90-0.12	16 57 46.5	-43 04 42	151	19.2(1.1)	117.1(1.5)	316.7(3.0)	419.0(6.9)	662	2100.0 (8.1)	N	8,24
G343.19-0.08(a)	16 58 34.9	-42 49 46	163	68.4(1.3)	146.0(0.7)	169.1(12.8)	193.9(19.8)	726	3732.1 (114.7)	N	24

Table 3—Continued

Name	J2000 Coordinates		area($''^2$)	Integrated flux(mJy)					area($''^2$)	Integrated flux(mJy)	Saturated?	Upper limit?
	α (h m s)	δ ($^\circ$ ' $''$)	IRAC	[3.6]	[4.5]	[5.8]	[8.0]	MIPS	[24]			
G343.40-0.40	17 00 40.4	-42 51 33	66	17.3(0.8)	34.5(0.8)	47.6(2.6)	40.2(7.5)	109	94.4 (10.1)		N	
G343.42-0.33	17 00 26.3	-42 48 03	127	36.8(0.2)	74.6(0.9)	122.4(0.9)	141.9(31.8)	167	298.7 (14.4)		N	
G343.50+0.03	16 59 10.7	-42 31 07	78	12.7(0.4)	27.4(0.5)	9.5(4.9)	31.8(1.3)	121	168.6 (13.9)		N	
G343.53-0.51(a)	17 01 32.2	-42 49 36	121	13.1(1.1)	62.4(0.3)	88.0(0.6)	75.0(10.2)	559	621.6 (17.0)		N	
G343.78-0.24	17 01 13.1	-42 27 48	204	23.3(2.1)	50.3(2.0)	39.7(7.9)	15.4(7.6)	766	2315.2 (36.2)		N	
G344.21-0.62	17 04 17.8	-42 21 09	200	111.5(0.2)	272.8(1.4)	358.0(4.3)	320.0(11.6)	645	1046.7 (351.3)		N	24
G348.17+0.46	17 12 10.9	-38 31 59	69	14.3(1.6)	44.4(1.2)	92.0(4.3)	145.9(9.8)	559	2432.5 (135.0)		N	24

Table 4. [4.5] Integrated Flux Densities: ”possible” MYSO outflow candidates

Name	J2000 Coordinates		area($''^2$)	Integrated flux(mJy)
	α (h m s)	δ ($^\circ$ ' ")	IRAC	[4.5]
G12.20-0.03	18 12 23.6	-18 22 54	192	228.9(0.2)
G12.42+0.50	18 10 51.1	-17 55 50	1767	1404.7(4.3)
G12.68-0.18	18 13 54.7	-18 01 47	300	159.8(4.6)
G17.96+0.08	18 23 21.0	-13 15 11	117	141.9(5.0)
G19.61-0.14	18 27 16.8	-11 53 51	53	21.7(1.3)
G20.24+0.07	18 27 44.6	-11 14 54	105	81.3(1.6)
G21.24+0.19	18 29 10.2	-10 18 11	20	6.3(0.3)
G23.82+0.38	18 33 19.5	-07 55 37	85	49.5(1.9)
G24.11-0.17	18 35 52.6	-07 55 17	147	592.6(1.0)
G24.11-0.18	18 35 53.0	-07 55 23	55	50.7(0.2)
G24.33+0.14	18 35 08.1	-07 35 04	104	67.8(2.7)
G28.85-0.23	18 44 47.5	-03 44 15	53	6.6(0.1)
G29.89-0.77	18 48 37.7	-03 03 44	45	7.9(0.4)
G29.91-0.81	18 48 47.6	-03 03 31	65	20.3(0.2)
G35.83-0.20	18 57 26.9	+02 29 00	42	3.1(0.5)
G39.39-0.14	19 03 45.3	+05 40 43	124	213.0(0.2)
G40.60-0.72	19 08 03.3	+06 29 15	120	73.7(1.6)
G43.04-0.45(a)	19 11 38.9	+08 46 39	52	19.5(0.2)
G43.04-0.45(b)	19 11 39.1	+08 46 32	27	9.6(0.1)
G45.47+0.13	19 14 07.3	+11 12 16	59	121.2(0.5)
G45.50+0.12	19 14 13.0	+11 13 30	30	6.3(0.1)
G49.91+0.37	19 21 47.5	+15 14 26	32	9.4(0.2)
G54.11-0.04	19 31 40.0	+18 43 53	29	11.7(0.1)
G54.11-0.05	19 31 42.2	+18 43 45	39	5.8(0.4)
G57.61+0.02	19 38 40.8	+21 49 35	42	9.0(0.5)
G58.78+0.64	19 38 49.6	+23 08 40	48	19.2(0.1)
G58.78+0.65	19 38 49.2	+23 08 50	33	6.2(0.2)
G305.77-0.25	13 16 30.0	-62 59 09	168	15.2(0.6)
G305.80-0.24	13 16 43.4	-62 58 29	22	5.8(0.1)
G309.38-0.13(b)	13 47 20.5	-62 18 11	27	7.5(0.5)
G309.97+0.50	13 51 05.2	-61 33 20	43	3.8(0.2)
G309.99+0.51(c)	13 51 12.9	-61 32 29	14	3.8(0.1)
G310.15+0.76	13 51 59.2	-61 15 37	516	324.4(0.9)
G310.38-0.30(a)	13 56 01.8	-62 14 15	40	4.0(0.1)
G310.38-0.30(b)	13 56 01.0	-62 14 19	46	3.0(0.5)
G310.38-0.30(c)	13 55 59.9	-62 14 27	52	1.5(0.1)
G310.38-0.30(d)	13 56 00.0	-62 14 18	29	1.3(0.2)
G311.04+0.69	13 59 18.1	-61 06 33	30	1.3(0.3)
G313.71-0.19(a)	14 22 37.4	-61 08 17	94	29.1(1.2)
G313.71-0.19(b)	14 22 35.6	-61 08 41	60	7.7(0.4)
G313.76-0.86	14 25 01.3	-61 44 57	210	156.7(0.9)
G317.44-0.37	14 51 03.0	-59 49 58	63	3.0(0.7)
G320.24-0.29	15 09 55.0	-58 25 31	29	8.4(0.7)
G323.74-0.26	15 31 45.5	-56 30 50	148	778.1(1.0)
G325.52+0.42	15 39 10.6	-54 55 40	23	7.0(0.1)

Table 4—Continued

Name	J2000 Coordinates		area($''^2$)	Integrated flux(mJy)
	α (h m s)	δ ($^\circ$ ' ")	IRAC	[4.5]
G326.37+0.94	15 41 44.1	-54 00 00	76	3.5(0.3)
G326.64+0.76	15 43 56.6	-53 58 32	39	3.3(0.2)
G326.99-0.03	15 49 07.8	-54 23 02	40	4.4(0.5)
G327.57-0.85	15 55 47.3	-54 39 09	125	373.9(2.7)
G327.65+0.13	15 52 00.5	-53 50 41	50	96.0(0.3)
G328.16+0.59	15 52 42.5	-53 09 51	251	53.0(0.7)
G328.60+0.27	15 56 15.8	-53 07 50	50	3.5(0.7)
G328.81+0.63	15 55 48.4	-52 43 06	851	1461.9(5.3)
G330.95-0.18	16 09 52.7	-51 54 56	181	225.1(0.7)
G331.12-0.46	16 11 55.3	-52 00 10	58	6.4(0.3)
G331.34-0.35	16 12 26.4	-51 46 17	94	460.4(1.6)
G331.51-0.34	16 13 11.7	-51 39 12	102	4.4(0.7)
G332.28-0.55	16 17 41.8	-51 16 04	53	8.2(0.2)
G332.29-0.09	16 15 45.2	-50 55 52	62	153.3(0.4)
G332.29-0.55	16 17 44.3	-51 15 42	26	3.2(0.7)
G332.35-0.44	16 17 31.4	-51 08 22	33	11.9(0.2)
G333.13-0.56	16 21 36.1	-50 40 49	386	38.7(4.2)
G333.32+0.10	16 19 28.9	-50 04 40	60	19.8(0.8)
G335.59-0.30	16 31 02.5	-48 44 07	176	21.8(0.6)
G336.02-0.83	16 35 09.7	-48 46 44	226	49.9(1.5)
G336.03-0.82	16 35 09.6	-48 45 55	35	10.8(0.3)
G337.40-0.40	16 38 50.4	-47 28 04	386	354.0(4.0)
G338.42-0.41	16 42 50.5	-46 42 29	256	30.1(2.5)
G338.92+0.55(a)	16 40 33.6	-45 41 44	104	125.8(0.6)
G338.92+0.55(b)	16 40 34.1	-45 42 07	230	750.6(2.3)
G339.58-0.13	16 45 59.5	-45 38 44	222	232.7(1.4)
G340.05-0.25	16 48 14.7	-45 21 52	128	27.6(2.7)
G340.06-0.23	16 48 09.7	-45 20 58	217	71.8(18.2)
G340.07-0.24	16 48 15.1	-45 20 57	23	2.6(0.1)
G340.10-0.18	16 48 07.0	-45 17 06	124	25.2(2.6)
G340.76-0.12	16 50 14.6	-44 44 38	30	3.7(0.4)
G341.22-0.26(b)	16 52 30.3	-44 28 40	109	24.9(2.3)
G342.04+0.43	16 52 27.8	-43 24 17	52	5.9(0.4)
G343.19-0.08(b)	16 58 35.8	-42 49 38	63	9.7(0.3)
G343.42-0.37	17 00 37.4	-42 49 40	17	1.9(0.1)
G343.53-0.51(b)	17 01 33.5	-42 49 50	52	13.6(0.1)
G344.22-0.57	17 04 06.4	-42 18 58	174	29.2(0.8)
G345.00-0.22(a)	17 05 11.2	-41 29 03	510	583.1(0.9)
G345.00-0.22(b)	17 05 10.9	-41 29 13	164	88.8(0.3)
G345.13-0.17	17 05 23.1	-41 21 11	78	34.4(0.2)
G346.04+0.05	17 07 19.9	-40 29 49	46	15.7(0.7)
G346.28+0.59	17 05 51.3	-39 58 43	268	324.4(1.5)
G348.18+0.48	17 12 08.0	-38 30 52	275	113.7(10.2)
G348.72-1.04	17 20 06.1	-38 57 15	36	33.0(5.0)
G348.73-1.04	17 20 06.5	-38 57 08	35	67.6(14.6)

Table 5. ”Outflow-only” [3.6] and [4.5] Integrated Flux Densities

Name	J2000 Coordinates		area($''^2$)	Integrated flux(mJy)		[3.6]-[4.5]
	α (h m s)	δ ($^{\circ}$ ' ")	IRAC	[3.6]	[4.5]	
G12.91-0.26	18 14 39.5	-17 52 00	965	172.3(13.2)	385.7(23.4)	0.87
G19.01-0.03 O-N ^a	18 25 44.8	-12 22 45.76	507	66.2(4.3)	141.3(3.0)	0.82
G19.01-0.03 O-S ^a	18 25 44.8	-12 22 45.76	98	11.7(0.7)	28.5(1.3)	0.97
G34.26+0.15	18 53 16.4	+01 15 07	3787	324.8(8.6)	838.9(15.6)	1.03
G37.55+0.20	18 59 07.5	+04 12 31	626	13.3(3.5)	39.2(5.4)	1.17
G345.51+0.35	17 04 24.6	-40 43 57	1407	244.4(11.4)	386.7(9.3)	0.50

^aCoordinates are for the central source of the EGO G19.01-0.03, as given in Table 1.

Table 6. Infrared EGO associations.

Name	IRDC?	Nearest IRAS PS ^a	Ang. Sep. (")
Table 1			
G11.92-0.61	Y	18110-1854	62
G12.02-0.21	Y	18097-1835	137
G12.91-0.03	Y	18111-1746	200
G14.33-0.64	Y	18159-1648	15
G14.63-0.58	Y	18164-1631	34
G16.61-0.24	Y	18190-1435	133
G18.67+0.03	N	18220-1241	36
G18.89-0.47	Y	18242-1241	113
G19.01-0.03	Y	18228-1224	88
G19.88-0.53	Y	18264-1152	6
G22.04+0.22	Y	18278-0936	5
G23.01-0.41	N	18318-0901	130
G23.96-0.11	N	18326-0802	113
G24.00-0.10	Y	18326-0802	50
G24.17-0.02	Y	18326-0751	64
G24.94+0.07	N	18337-0707	78
G25.27-0.43	Y	18362-0703	29
G27.97-0.47	Y	18412-0440	116
G28.83-0.25	Y	18421-0348	35
G34.41+0.24	Y	18507+0121	31
G35.03+0.35	N	18515+0157	58
G35.04-0.47	Y	18545+0134	135
G35.13-0.74	N	18553+0133	215
G35.15+0.80	N	18500+0216	24
G35.20-0.74	N	18556+0136	4
G35.68-0.18	Y	18547+0215	246
G35.79-0.17	Y	18547+0223	10
G36.01-0.20	Y	18551+0238	194
G37.48-0.10	N	18574+0355	191
G39.10+0.49	N	18585+0538	5
G40.28-0.27	Y	19034+0618	70
G44.01-0.03	N	19094+0944	79
G45.47+0.05	Y	19120+1103	59
G49.27-0.34	Y	19207+1410	221
G54.45+1.02	N	19262+1925	12
G56.13+0.22	N	19326+2030	19
G58.09-0.34	Y	19388+2156	23
G59.79+0.63	Y	19389+2354	10
G298.26+0.74	N	12091-6129	3
G298.89+0.37	Y	12140-6157	76
G298.90+0.36	N	12140-6157	19
G305.48-0.10	N	13108-6233	191
G305.52+0.76	Y	13102-6143	12
G305.57-0.34	Y	13113-6249	109
G305.62-0.34	N	13119-6249	7

Table 6—Continued

Name	IRDC?	Nearest IRAS PS ^a	Ang. Sep. (")
G305.82-0.11	Y	13128-6236	291
G305.89+0.02	Y	13140-6226	27
G309.15-0.35	Y	13421-6217	125
G309.38-0.13(a)	Y	13438-6203	32
G309.99+0.51(a)	Y	13475-6115	129
G310.08-0.23	Y	13497-6159	31
G312.11+0.26	Y	14050-6056	171
G317.42-0.67	Y	14481-5953	5
G317.46-0.40(b)	Y	14480-5941	313
G317.87-0.15	Y	14488-5915	276
G317.88-0.25	Y	14500-5918	78
G321.94-0.01	Y	15158-5707	34
G324.19+0.41	N	15277-5532	39
G324.72+0.34	Y	15310-5517	13
G326.27-0.49	Y	15432-5501	7
G326.31+0.90	Y	15375-5352	168
G326.78-0.24	N	15450-5431	7
G326.79+0.38	Y	15425-5401	2
G326.86-0.67	Y	15475-5449	104
G326.97-0.03	Y	15453-5416	164
G327.12+0.51	N	15437-5343	4
G327.39+0.20	N	15464-5348*	4
G327.40+0.44	Y	15454-5335	29
G328.14-0.43	Y	15529-5350	94
G329.18-0.31	Y	15579-5303	8
G329.47+0.52	Y	15557-5215	53
G329.61+0.11	Y	15584-5230	208
G332.35-0.12	Y	16122-5047	18
G332.56-0.15	Y	16132-5039	97
G332.81-0.70	N	16170-5053	48
G332.94-0.69	Y	16175-5046	22
G332.96-0.68	Y	16175-5045	3
G333.18-0.09	Y	16159-5012	75
G335.59-0.29	Y	16272-4837	21
G337.30-0.87	Y	16366-4746	82
G338.39-0.40	Y	16390-4637	25
G339.95-0.54	N	16455-4531	90
G340.97-1.02	Y	16513-4504	48
G341.24-0.27	Y	16487-4423	170
G341.73-0.97	Y	16537-4426	3
G341.99-0.10	Y	16510-4341	55
G343.12-0.06	Y	16547-4247	4
G343.50-0.47	N	16579-4245	120
G343.72-0.18(a)	Y	16572-4221	137
G343.72-0.18(b)	Y	16574-4225	133
G344.58-0.02	N	16594-4137	9

Table 6—Continued

Name	IRDC?	Nearest IRAS PS ^a	Ang. Sep.('')
G345.72+0.82	Y	16596-4012	0.4
G345.99-0.02	N	17039-4030	14
G347.08-0.40	Y	17089-3951	9
G348.55-0.98	N	17158-3901	64
G348.58-0.92	Y	17157-3855	129
G349.15-0.98	Y	17176-3831	0.8
	70/97	avg./median	75/50
Table 2			
G10.29-0.13	Y	18060-2005	134
G10.34-0.14	Y	18060-2005	111
G16.59-0.05	Y	18182-1433	19
G19.36-0.03	Y	18236-1205	24
G19.61-0.12	N	18244-1155	43
G25.38-0.15	Y	18355-0650	110
G28.28-0.36	N	18416-0420	81
G34.39+0.22	Y	18507+0121	52
G48.66-0.30	Y	19199+1347	463
G49.42+0.33	N	19186+1440	34
G309.90+0.23	Y	13475-6135	54
G317.46-0.40(a)	Y	14480-5941	319
G318.04+0.09	Y	14498-5856	24
G318.05+0.09	N	14498-5856	9
G320.23-0.28	Y	15061-5814	82
G324.17+0.44	N	15277-5532	102
G326.32-0.39	N	15428-5453	213
G326.47+0.70	Y	15394-5358	27
G326.48+0.70	Y	15394-5358	20
G326.61+0.80(b)	Y	15395-5348	89
G328.25-0.53	N	15541-5349	3
G329.03-0.20	Y	15566-5304	2
G329.41-0.46	N	15596-5301	11
G329.47+0.50	Y	15557-5215	48
G330.88-0.37	Y	16065-5158	20
G331.13-0.24	N	16071-5142	28
G332.12+0.94	Y	16067-5010	10
G332.60-0.17	Y	16136-5038	69
G332.73-0.62	Y	16158-5055	262
G333.47-0.16	N	16175-5002	14
G335.06-0.43	Y	16256-4905	17
G335.79+0.18	Y	16259-4805	224
G337.91-0.48	N	16374-4701	42
G341.22-0.26(a)	Y	16487-4423	121
G342.48+0.18	Y	16515-4308	51
G344.23-0.57	Y	17006-4215	94
	25/36	avg./median	84/50
Table 3			

Table 6—Continued

Name	IRDC?	Nearest IRAS PS ^a	Ang. Sep. (")
G11.11-0.11	Y	18073-1923	104
G16.58-0.08	Y	18182-1433	125
G24.63+0.15	Y	18331-0717	281
G29.84-0.47	Y	18446-0303	190
G29.96-0.79	Y	18461-0304	35
G34.28+0.18	Y	18507+0110	170
G40.28-0.22	Y	19031+0621	73
G45.80-0.36	N	19141+1110	10
G49.07-0.33	Y	19205+1403	179
G49.27-0.32	N	19207+1410	252
G50.36-0.42	Y	19234+1510	144
G53.92-0.07	N	19291+1826	20
G54.11-0.08	N	19294+1836	89
G54.45+1.01	N	19262+1925	9
G58.79+0.63	N	19366+2301	105
G62.70-0.51	N	19497+2549	14
G304.89+0.64	Y	13050-6154	6
G309.91+0.32	Y	13475-6129	69
G309.97+0.59	Y	13475-6115	169
G309.99+0.51(b)	Y	13475-6115	142
G311.51-0.45	Y	14023-6150A	101
G324.11+0.44	Y	15272-5533	2
G326.36+0.88	Y	15375-5352	308
G326.41+0.93	Y	15384-5348	160
G326.57+0.20	N	15422-5418	112
G326.61+0.80(a)	Y	15395-5348	102
G326.61+0.80(c)	Y	15395-5348	95
G326.65+0.75	Y	15402-5349	12
G326.80+0.51	Y	15415-5356	217
G326.92-0.31	Y	15462-5428	110
G327.30-0.58	Y	15492-5426	101
G327.72-0.38	N	15508-5403	84
G327.86+0.19	N	15489-5333	121
G327.89+0.15	Y	15489-5333	228
G328.55+0.27	Y	15522-5301	12
G329.07-0.31(a)	Y	15573-5307	17
G329.07-0.31(b)	Y	15573-5307	1
G329.16-0.29	Y	15579-5303	120
G329.18-0.30	Y	15579-5303	30
G331.08-0.47	Y	16076-5154	161
G331.37-0.40	Y	16089-5137	154
G331.62+0.53	N	16061-5048	21
G331.71+0.58	Y	16062-5041	70
G331.71+0.60	Y	16062-5041	3
G332.28-0.07	Y	16119-5048	100
G332.33-0.12	Y	16122-5047	61

Table 6—Continued

Name	IRDC?	Nearest IRAS PS ^a	Ang. Sep. (")
G332.36+0.60	Y	16093-5015	38
G332.47-0.52	Y	16147-5100	47
G332.58+0.15	Y	16122-5028	167
G332.59+0.04(a)	Y	16128-5033	185
G332.59+0.04(b)	Y	16128-5033	193
G332.91-0.55	Y	16168-5044	156
G333.08-0.56	Y	16172-5032	278
G334.04+0.35	N	16178-4916	7
G334.25+0.07	Y	16200-4919	48
G335.43-0.24	Y	16264-4841	77
G336.87+0.29	Y	16301-4718	138
G336.96-0.98	Y	16358-4804	32
G337.16-0.39	Y	16340-4732	22
G338.32-0.41	Y	16389-4639	129
G340.75-1.00	Y	16506-4512	161
G340.77-0.12	Y	16465-4437	126
G340.78-0.10	N	16465-4437	52
G341.20-0.26	Y	16487-4423	110
G341.23-0.27	Y	16487-4423	141
G342.15+0.51	N	16489-4318	453
G342.90-0.12	N	16541-4259	41
G343.19-0.08(a)	Y	16550-4245	10
G343.40-0.40	Y	16566-4249	345
G343.42-0.33	Y	16573-4240	342
G343.50+0.03	Y	16558-4228	158
G343.53-0.51(a)	Y	16579-4245	45
G343.78-0.24	Y	16576-4223	8
G344.21-0.62	N	17006-4215	95
G348.17+0.46	N	17089-3834	383
	58/75	avg./median	116/102
Table 4			
G12.2-0.03	N	18094-1823	22
G12.42+0.50	N	18079-1756	12
G12.68-0.18	N	18112-1801	206
G17.96+0.08	N	18205-1316	12
G19.61-0.14	N	18244-1155	13
G20.24+0.07	N	18249-1116	10
G21.24+0.19	Y	18263-1020	19
G23.82+0.38	N	18305-0758	22
G24.11-0.17	Y	18334-0757	216
G24.11-0.18	Y	18334-0757	210
G24.33+0.14	Y	18324-0737	5
G28.85-0.23	N	18421-0348	78
G29.89-0.77	Y	18460-0307	25
G29.91-0.81	N	18460-0307	128
G35.83-0.20	Y	18549+0226	83

Table 6—Continued

Name	IRDC?	Nearest IRAS PS ^a	Ang. Sep. (″)
G39.39-0.14	N	19012+0536	4
G40.60-0.72	N	19056+0624	8
G43.04-0.45(a)	N	19092+0841	24
G43.04-0.45(b)	N	19092+0841	25
G45.47+0.13	N	19117+1107	21
G45.50+0.12	N	19117+1107	91
G49.91+0.37	N	19195+1508	5
G54.11-0.04	N	19294+1836	89
G54.11-0.05	N	19294+1836	73
G57.61+0.02	N	19365+2142	18
G58.78+0.64	N	19366+2301	23
G58.78+0.65	N	19366+2301	21
G305.77-0.25	N	13134-6242	91
G305.80-0.24	N	13134-6242	17
G309.38-0.13(b)	Y	13438-6203	32
G309.97+0.50	Y	13475-6115	184
G309.99+0.51(c)	Y	13475-6115	149
G310.15+0.76	N	13484-6100	14
G310.38-0.30(a)	Y	13524-6159	9
G310.38-0.30(b)	Y	13524-6159	13
G310.38-0.30(c)	Y	13524-6159	24
G310.38-0.30(d)	Y	13524-6159	17
G311.04+0.69	Y	13558-6051	57
G313.71-0.19(a)	Y	14188-6054	37
G313.71-0.19(b)	Y	14188-6054	33
G313.76-0.86	Y	14212-6131	11
G317.44-0.37	Y	14467-5939	242
G320.24-0.29	Y	15061-5814	64
G323.74-0.26	N	15278-5620	30
G325.52+0.42	N	15353-5445	12
G326.37+0.94	Y	15375-5352	233
G326.64+0.76	Y	15402-5349	52
G326.99-0.03	Y	15453-5416	151
G327.57-0.85	Y	15519-5430	29
G327.65+0.13	N	15481-5341	2
G328.16+0.59	N	15488-5300	1
G328.60+0.27	Y	15522-5301	163
G328.81+0.63	Y	15520-5234	4
G330.95-0.18	Y	16060-5146	35
G331.12-0.46	Y	16082-5150	158
G331.34-0.35	N	16085-5138	41
G331.51-0.34	Y	16093-5128	168
G332.28-0.55	Y	16141-5107	159
G332.29-0.09	N	16119-5048	10
G332.29-0.55	Y	16141-5107	129
G332.35-0.44	Y	16137-5100	10

Table 6—Continued

Name	IRDC?	Nearest IRAS PS ^a	Ang. Sep.($''$)
G333.13-0.56	N	16178-5029	237
G333.32+0.10	Y	16157-4957	2
G335.59-0.30	Y	16272-4837	63
G336.02-0.83	N	16313-4840	33
G336.03-0.82	N	16313-4840	27
G337.40-0.40	Y	16351-4722	8
G338.42-0.41	Y	16390-4637	95
G338.92+0.55(a)	N	16368-4538	171
G338.92+0.55(b)	N	16368-4538	151
G339.58-0.13	Y	16424-4531	139
G340.05-0.25	Y	16445-4516	34
G340.06-0.23	Y	16445-4516	43
G340.07-0.24	Y	16445-4516	51
G340.10-0.18	N	16442-4511	169
G340.76-0.12	Y	16465-4437	130
G341.22-0.26(b)	Y	16487-4423	102
G342.04+0.43	N	16489-4318	77
G343.19-0.08(b)	Y	16550-4245	22
G343.42-0.37	Y	16573-4240	331
G343.53-0.51(b)	Y	16579-4245	51
G344.22-0.57	Y	17006-4215	89
G345.00-0.22(a)	Y	17016-4124	16
G345.00-0.22(b)	Y	17016-4124	15
G345.13-0.17	N	17020-4117	129
G346.04+0.05	N	17038-4026	60
G346.28+0.59	N	17023-3954	16
G348.18+0.48	N	17089-3820	404
G348.72-1.04	Y	17167-3854	40
G348.73-1.04	Y	17167-3854	39
	49/90	avg./median	73/36
Outflow-only (Table 5)			
G12.91-0.26	N	18117-1753	8
G34.26+0.15	N	18507+0110	56
G37.55+0.20	N	18566+0408	39
G345.51+0.35	N	17008-4040	55
	0/4	avg./median	40/47

^aB1950 coordinates

Table 7. 6.7GHz CH₃OH maser associations of EGOs

Name	Maser Survey ^a	Walsh IRAS target ^b	Maser? ^a	Offset(") ^c	Distance(kpc) ^d
Table 1					
G11.92-0.61	W	18110-1854	N		
G14.33-0.64	W	18159-1648	N		
G28.83-0.25	W	18421-0348	Y	3	
G298.26+0.74	W	12091-6129	Y	<1	
G312.11+0.26	W	14050-6056	Y	1	
G326.27-0.49	E		N		
G326.78-0.24	E		N		
G326.79+0.38	E		N		
G326.86-0.67 ^e	E		Y	5	3.7
G326.97-0.03	E		N		
G327.12+0.51	W,E	15437-5343	Y (W,E)	1	5.5
G327.39+0.20	E		Y	<1	5.3
G327.40+0.44	E		Y	5	5.2
G328.14-0.43	E		N		
G329.18-0.31	E		Y	3	3.6
G329.47+0.52	W,E	15557-5215	Y (W)	<1	4.5
G329.61+0.11	E		Y	<1	3.9
G332.35-0.12	C,E		N		
G332.56-0.15	C,W,E	16132-5039	Y (C,W,E)	2	3.5
G332.94-0.69 ^e	E		Y	1	3.6
G333.18-0.09	C,E	16159-5012	Y(C,E, W could not locate)	1	5
G335.59-0.29	C,W,E	16272-4837	Y (C,W)	2,4,17	
G339.95-0.54	W	16455-4531	Y	1	
G343.12-0.06	W	16547-4247	N		
G344.58-0.02	W	16594-4137	Y	<1	
Table 2					
G10.29-0.13	W	18060-2005	Y	2	
G10.34-0.14	W	18060-2005	Y	<1	
G16.59-0.05	W	18182-1433	Y	<1	
G19.36-0.03	W	18236-1205	Y	4	
G19.61-0.12	W	18244-1155	Y	5	
G28.28-0.36	W	18416-0420	Y	1	
G320.23-0.28	W	15061-5814	Y	6	
G326.32-0.39	E		N		
G326.47+0.70 ^e	E		Y?	11	2.6
G326.48+0.70 ^e	E		Y?	8	2.6
G328.25-0.53	E		Y	1	2.6
G329.03-0.20	E		Y	7,19	2.6,2.9
G329.41-0.46	W,E	15596-5301	Y(E)	3	4.2
G329.47+0.50	W,E	15557-5215	Y	3	4.5
G330.88-0.37	W,E	16065-5158	N		
G331.13-0.24	W,E	16071-5142	Y (W,E)	3	5.2
G332.60-0.17	C,E		Y(C,E)	<1	3.5
G332.73-0.62	W	16158-5055	Y	1	
G333.47-0.16	C,W,E	16175-5002	Y (C,W,E)	1	3

Table 7—Continued

Name	Maser Survey ^a	Walsh IRAS target ^b	Maser? ^a	Offset(") ^c	Distance(kpc) ^d
G335.06-0.43	E		Y	<1	3.4
G335.79+0.18	C		Y	5	
G337.91-0.48	W	16374-4701	N		
G344.23-0.57	W	17006-4215	Y	7,8	
Table 3					
G16.58-0.08	W	18182-1433	N		
G326.57+0.20	E		N		
G326.80+0.51	E		N		
G326.92-0.31	E		N		
G327.72-0.38	E		N		
G327.86+0.19	E		N		
G327.89+0.15	E		N		
G328.55+0.27	E		N		
G329.07-0.31(a)	E		N		
G329.07-0.31(b)	E		Y	<1	3
G329.16-0.29	E		N		
G329.18-0.30	E		N		
G331.08-0.47	E		N		
G331.37-0.40	W,E	16085-5138	N		
G331.62+0.53	E		N		
G332.28-0.07	C,E		N		
G332.33-0.12	C,E		N		
G332.47-0.52	E		N		
G332.58+0.15	C,E		N		
G332.59+0.04(a)	C,E		N		
G332.59+0.04(b)	C,E		N		
G334.04+0.35	E		N		
G334.25+0.07	C,E		N		
G340.75-1.00	W	16506-4512	N		
G343.19-0.08(a)	W	16547-4247	N		
G344.21-0.62	W	17006-4215	N		
Table 4					
G19.61-0.14	W	18244-1155	N		
G28.85-0.23	W	18421-0348	Y	2	
G305.77-0.25	W	13134-6242	N		
G305.80-0.24	W	13134-6242	N		
G309.97+0.50	W	13471-6120	N		
G313.76-0.86	W	14212-6131	Y	3,25	
G320.24-0.29	W	15061-5814	N		
G323.74-0.26	W	15278-5620	Y	0.6,0.8,3	
G325.52+0.42	E		N		
G326.99-0.03	E		N		
G327.65+0.13	E		N		
G328.60+0.27	E		N		
G328.81+0.63	W,E	15520-5234	Y (W,E)	<1	3
G330.95-0.18	C,W,E	16060-5146	Y (W,E)	3	5.3

Table 7—Continued

Name	Maser Survey ^a	Walsh IRAS target ^b	Maser? ^a	Offset(") ^c	Distance(kpc) ^d
G331.12-0.46	E		N		
G331.34-0.35	W,E	16085-5138	Y(W,E)	1	4.3
G331.51-0.34	E		N		
G332.29-0.09	C,E		Y(E)	3	3.2
G332.35-0.44	E		Y	2	3.6
G333.13-0.56 ^e	E		Y	10	3.8
G333.32+0.10	C,E		Y(C,E)	2	3.1
G335.59-0.30	W	16272-4837	N		
G336.02-0.83	W	16313-4840	Y	5	
G336.03-0.82	W	16313-4840	N		
G337.40-0.40	W	16351-4722	Y	3,4	
G339.58-0.13	C	16424-4531 and 16421-4532	Y	8	
G340.05-0.25	W	16445-4516	Y	12	
G340.06-0.23	W	16445-4516	N		
G340.07-0.24	W	16445-4516	N		
G343.19-0.08(b)	W	16547-4247	N		
G344.22-0.57	W	17006-4215	N		
G345.00-0.22(a)	W	17016-4124	Y	4,5	
G345.00-0.22(b)	W	17016-4124	N		
G348.72-1.04	W	17167-3854	N		
G348.73-1.04	W	17167-3854	Y	<1	
Outflow-only (Table 5)					
G12.91-0.26	W	18117-1753	Y	<1	
G345.51+0.35	W	17008-4040	Y	32	

^aC=Caswell (1996), W=Walsh et al. (1998), E=Ellingsen (2006)

^bB1950 coordinates

^cNominal separation from EGO positions listed in Tables 1-5, rounded to the nearest arcsecond. Separations less than 1" are denoted <1. Where multiple offsets are listed, multiple spatially distinct maser spots are reported.

^dDistances are the near kinematic distances reported in Ellingsen (2006). The association of EGOs with IRDCs supports the adoption of the near kinematic distance.

^eSource outside the nominal area of Ellingsen et al. (1996) and Ellingsen (2006) of $325^\circ < l < 335^\circ$, $b=\pm 0.53^\circ$ and not included in the calculation of the fraction of EGOs within the survey coverage that have associated 6.7 GHz CH₃OH masers.




Fig. 1.— (a-e): IRAC (a) $3.6\ \mu\text{m}$, (b) $4.5\ \mu\text{m}$, (c) $5.8\ \mu\text{m}$, (d) $8.0\ \mu\text{m}$ and (e) MIPS $24\ \mu\text{m}$ images of the EGO G11.92-0.61. (f): Three-color GLIMPSE IRAC image of the EGO G11.92-0.61 showing $8.0\ \mu\text{m}$ (red), $4.5\ \mu\text{m}$ (green), and $3.6\ \mu\text{m}$ (blue). The polygonal source aperture used for IRAC photometry is overlaid in all panels. The circular region used for MIPS photometry is overlaid on the $24\ \mu\text{m}$ image. All panels are on the same spatial scale. This is an example of a source assigned to Table 1.




Fig. 2.— (a) Three-color GLIMPSE IRAC image of the EGO G11.92-0.61 showing $8.0\ \mu\text{m}$ (red), $4.5\ \mu\text{m}$ (green), and $3.6\ \mu\text{m}$ (blue), with the 3 polygonal apertures chosen for IRAC photometry by three different observers overlaid. (b) MIPS $24\ \mu\text{m}$ image of the EGO G11.92-0.61 with the 3 circular apertures chosen for MIPS photometry by three different observers overlaid. The MIPS counterpart is slightly saturated, indicated by the dip in flux at its center.




Fig. 3.— Same as Figure 1, but for the EGO G10.29-0.13. The polygonal source aperture used for IRAC photometry is overlaid in all panels. This is an example of a source assigned to Table 2 because of the lack of clear counterparts in bands other than [4.5].




Fig. 4.— Same as Figure 1, but for the EGO G10.34-0.14, an example of a source assigned to Table 2 because of the lack of a clear counterpart at $8.0\ \mu\text{m}$ and confusion in the $24\ \mu\text{m}$ image. The polygonal source aperture used for IRAC photometry is overlaid in all panels.




Fig. 5.— Same as Figure 1, but for the EGO G11.11-0.11, an example of a source assigned to Table 3 because, while clear counterparts are visible in all bands, the green emission is not very extended and surrounds a bright multiband IRAC source. The polygonal source aperture used for IRAC photometry is overlaid in all panels. The circular region used for MIPS photometry is overlaid on the 24 μm image.




Fig. 6.— Same as Figure 1, but for the EGO G12.20-0.03, an example of a source assigned to Table 4 because of the limited extent of the $4.5\ \mu\text{m}$ emission and the degree of confusion in the region. At least 2 multiband IRAC sources are apparent, either or both of which could be associated with the extended $4.5\ \mu\text{m}$ emission; $24\ \mu\text{m}$ emission from several IRAC sources appears to be blended in a single, saturated, MIPS counterpart. The polygonal source aperture used for IRAC photometry is overlaid in all panels.




Fig. 7.— Histogram, binned to $0.2'$, of number of EGOs as a function of angular separation from the nearest *IRAS* point source. About $\sim 50\%$ of the catalogued EGO population is $>1'$ from the nearest *IRAS* point source.




Fig. 8.— The EGO G11.92-0.61 as seen by (a) *IRAS* (red: 100 μm , green: 25 μm , blue: 12 μm), (b) *MSX* (red: 21.4 μm , green: 12.1 μm , blue: 8.3 μm), and (c) *Spitzer* IRAC (red: 8.0 μm , green: 4.5 μm , blue: 3.6 μm). Error ellipses for the two IRAS point sources in the field are overplotted. All panels are on the same spatial scale.




Fig. 9.— Locations of EGOs overplotted on contours of integrated CO emission from Dame et al. (2001). EGOs in Table 1 are shown as diamonds, Table 2 as squares, Table 3 as asterisks, Table 4 as triangles, and Table 5 as crosses. CO contour levels are 39.5, 79.0, 118.6, 158.1, 197.6, 237.2 K km s⁻¹. The overall distribution pattern of EGOs with respect to CO is the same for sources from all subcatalogs: the vast majority of EGOs coincide with CO clouds.




Fig. 10.— Same as Figure 3, but with a 6.7 GHz CH_3OH maser from Walsh et al. (1998) marked. The region used for IRAC photometry is also overplotted. Similar images are available online for all EGOs associated with 6.7 GHz CH_3OH masers in the surveys of Caswell (1996), Walsh et al. (1998) and Ellingsen (2006).




Fig. 11.— Histogram, binned to $1.0''$, of number of EGOs as a function of angular separation from associated 6.7 GHz CH_3OH masers. This nominal offset is small ($\lesssim 5''$ for the majority of EGOs). This is consistent with our criteria for including an EGO in Table 7: a maser spatially coincident either with extended $4.5\ \mu\text{m}$ emission or with a $24\ \mu\text{m}$ or multiband IRAC source likely to be associated with extended $4.5\ \mu\text{m}$ emission.




Fig. 12.— (a-e): IRAC (a) $3.6\ \mu\text{m}$, (b) $4.5\ \mu\text{m}$, (c) $5.8\ \mu\text{m}$, (d) $8.0\ \mu\text{m}$ and (e) MIPS $24\ \mu\text{m}$ images of the EGO G19.01-0.03. The polygonal aperture used for the photometry reported in Table 1 is shown in all panels; the aperture used for MIPS $24\ \mu\text{m}$ photometry is also shown in panel (e). (f): Three-color GLIMPSE IRAC image of the EGO G19.01-0.03 showing $8.0\ \mu\text{m}$ (red), $4.5\ \mu\text{m}$ (green), and $3.6\ \mu\text{m}$ (blue) with the polygonal source apertures used for “outflow-only” IRAC photometry reported in Table 5 overlaid. All panels are on the same spatial scale. The central point source dominates the flux density in the IRAC Table 1 aperture at wavelengths longer than $4.5\ \mu\text{m}$.




Fig. 13.— Color-color plot of EGOs in Table 1 (filled circles) and Table 3 (open circles). The error bar at top left was derived from the average standard deviation of the measurements (Tables 1 and 3), converted to magnitudes and propagated to colors. The EGO points are overlaid on a grid of YSO models in greyscale (Robitaille et al. 2006). The black lines indicate regions occupied predominantly by models of various evolutionary stages: at right are the youngest sources surrounded by infalling envelopes (Stage I); in the middle are more evolved sources surrounded by disks (Stage II), and at left are sources with low-mass disks (Stage III). Note that most EGOS fall in the region occupied by the youngest protostar models. EGOs with fluxes flagged as saturated or as upper limits in Tables 1 and 3 were excluded from the plot.